

Problem selection

Vladimir Barzov

This is a collection of original math problems and their solutions. The problems have been categorized as per IMO's standard into algebra, geometry, combinatorics, and number theory sections.

1 Algebra

1.1

Let $n > 1$.

$$0 = a_1 < a_2 < \dots < a_n, \quad 0 = b_n < b_{n-1} < \dots < b_1.$$

We set

$$\alpha_i = \sqrt{1 - \sqrt{\frac{1}{1 + \left(\frac{a_i b_{i+1} - b_i a_{i+1}}{a_i a_{i+1} + b_i b_{i+1}}\right)^2}}}, \quad i = 1, 2, \dots, n-1.$$

Prove that

$$\sum_{i=1}^{n-1} \alpha_i < 1.111.$$

Solution: From $a_i^2 + b_i^2 > 0$, without loss of generality we may replace $\{a_i, b_i\}$ by

$$\left\{ \frac{a_i}{\sqrt{a_i^2 + b_i^2}}, \frac{b_i}{\sqrt{a_i^2 + b_i^2}} \right\}, \quad i = 1, 2, \dots, n,$$

without changing the values of $\alpha_1, \alpha_2, \dots, \alpha_{n-1}$ as the expression inside the innermost brackets is homogeneous in a_i and b_i , hence we can scale the a_i and b_i by the same factor to get $a_i^2 + b_i^2 = 1$. Now, the expression for α_i simplifies to:

$$\begin{aligned} \alpha_i &= \sqrt{1 - \frac{a_i a_{i+1} + b_i b_{i+1}}{\sqrt{(a_i^2 + b_i^2)(a_{i+1}^2 + b_{i+1}^2)}}} = \sqrt{1 - (a_i a_{i+1} + b_i b_{i+1})} \\ &= \frac{1}{\sqrt{2}} \sqrt{1 - 2a_i a_{i+1} - 2b_i b_{i+1}} = \frac{1}{\sqrt{2}} \sqrt{(a_i - a_{i+1})^2 + (b_i - b_{i+1})^2}. \end{aligned}$$

Consider a coordinate system and points with coordinates (a_i, b_i) . They lie in the first quadrant of the circle $x^2 + y^2 = 1$. Since a_i increases and b_i decreases, the ratios a_i/b_i strictly


increase, hence the angles $\angle A_i O x$ strictly decrease. Therefore the points are distinct and appear in clockwise order in the first quadrant, with A_1 lying exactly on the y -axis. Then

$$\sqrt{2} \sum_{i=1}^{n-1} \alpha_i$$

represents the length of the polygonal line $A_1 A_2 \cdots A_n$. Since each $|A_i A_{i+1}|$ is a chord in a circle and is strictly shorter than the arc $\widehat{A_i A_{i+1}}$, the length of this line is strictly shorter than the sum of the lengths of the arcs, which in turn is no greater than a quarter of the perimeter of the circle, namely $\pi/2$. It follows that

$$\sum_{i=1}^{n-1} \alpha_i < \frac{1}{\sqrt{2}} \cdot \frac{\pi}{2} = 1.1107 < 1.111.$$

Notice that the bound is quite sharp. □



1.2

A sequence of pairs of numbers $(a_1, b_1), (a_2, b_2), \dots, (a_{2001}, b_{2001})$ is given such that

$$(a_1, b_1) = (0, 1)$$

and for $n = 1, 2, \dots, 2000$ either

$$(a_{n+1}, b_{n+1}) = (-a_n, b_n)$$

or

$$(a_{n+1}, b_{n+1}) = \left(\frac{5}{4}a_n + \frac{3}{4}b_n, \frac{5}{4}b_n + \frac{3}{4}a_n \right).$$

Prove that

$$(a_1 + b_1)(a_2 + b_2), \dots, (a_{2001} + b_{2001}) \geq 1.$$

Solution: Let $\varepsilon = \ln 2$, so that $\sinh(\varepsilon) = 3/4$ and $\cosh(\varepsilon) = 5/4$. We claim that the representation

$$(a_n, b_n) = (\sinh(\alpha_n), \cosh(\alpha_n)) \quad (*)$$

holds for $\alpha_1 = 0$. Next, for $n \geq 1$, either

$$(a_{n+1}, b_{n+1}) = (-a_n, b_n) = (\sinh(-\alpha_n), \cosh(\alpha_n))$$

or

$$\begin{aligned} (a_{n+1}, b_{n+1}) &= \left(\frac{5}{4}a_n + \frac{3}{4}b_n, \frac{5}{4}b_n + \frac{3}{4}a_n \right) \\ &= (\cosh \varepsilon \sinh \alpha_n + \sinh \varepsilon \cosh \alpha_n, \cosh \varepsilon \cosh \alpha_n + \sinh \varepsilon \sinh \alpha_n) \\ &= (\sinh(\alpha_n + \varepsilon), \cosh(\alpha_n + \varepsilon)). \end{aligned}$$

Therefore, the identity $(*)$ remains valid for all n where

$$\alpha_{n+1} = -\alpha_n \quad \text{or} \quad \alpha_{n+1} = \alpha_n + \varepsilon.$$

We shall prove the following

Lemma.

$$\alpha_1 + \alpha_2 + \dots + \alpha_n \geq 0 \quad \text{for } n \geq 1.$$

Proof: Let us encode the given sequence satisfying the above properties by means of a symbolic string of x and y (whose length is one less than the length of the sequence), corresponding to the case $\alpha_{n+1} = -\alpha_n$ or $\alpha_{n+1} = \alpha_n + \varepsilon$. For example, for the sequence

$$0, \varepsilon, 2\varepsilon, -2\varepsilon, -\varepsilon, \varepsilon$$

the symbolic string representation is $yyxyx$. For completeness, let the empty string correspond to the single-term sequence α_1 . For every such string A we define by $s(A)$ the sum of the members of the sequence. We shall make the following observations:

- Observation 1. For every symbolic sequence A ,

$$s(A) = s(Axx) = s(Ayx),$$

because if α is the last term of the sequence encoded by A , the only difference with Axx is the last two terms $(-\alpha), \alpha$, and the only difference with Ayx is the last two terms $\alpha + \varepsilon, -(\alpha + \varepsilon)$. In both cases the terms sum to 0.

- Observation 2. A leading x replaces $\alpha_1 = 0$ with the pair $(0, 0)$. Therefore any number of leading x 's simply inserts zeros into the sequence and does not change the sum, hence

$$s(xA) = s(A).$$

- Observation 3. If α is the last term of the sequence encoded by A , the only difference between $AyxyB$ and AxB is the two terms following A , namely $\alpha + \varepsilon, -(\alpha + \varepsilon)$, after which point the two sequences continue with $-\alpha$ the same way. Therefore

$$s(AyxyB) = s(AxB).$$

To prove that there is no sequence with negative sum, assume the contrary, and let's pick the shortest such symbolic string. We can assume it has more than 1 symbols, as the cases for 0 and 1 can easily be examined: the empty string gives rise to 0, and the one-symbol string sequences x and y give sums 0 and ε respectively. In any of these cases, the sums are non-negative. So the shortest sequence must have more than 1 symbols.

By the symbolic analogue of this sequence, the string does not begin with x (due to observation 2), as it could otherwise be reduced to a shorter one that still has negative sum. Furthermore, it does not end with x either due to observation 1, as we could otherwise chop the last two symbols regardless if they are xx or yx . Then it must begin and end with y , and according to observation 3, it must not contain x , else it would be reducible to a shorter one. So the string contains only y , and hence its corresponding α sequence is

$$0, \varepsilon, 2\varepsilon, \dots, k\varepsilon$$

which has nonnegative sum – a contradiction. Thus the lemma is proved and so

$$\prod_{i=1}^{2001} (a_i + b_i) = \prod_{i=1}^{2001} (\sinh(\alpha_i) + \cosh(\alpha_i)) = \prod_{i=1}^{2001} e^{\alpha_i} = e^{\sum_{i=1}^{2001} \alpha_i} \geq 1.$$

1.3

Given a finite set A_0 of *rational* numbers. We define the sequence of sets

$$A_{k+1} = \left\{ \frac{x+y}{1+xy} \mid x, y \in A_k, xy \neq -1 \right\}, \quad k = 0, 1, \dots$$

Prove that there exists a *rational* number which does not belong to any of the sets A_k , for $k = 0, 1, \dots$

Solution: If 1 is not in A_0 , then it is not in any of the subsequent sets since

$$\frac{x+y}{1+xy} = 1 \Rightarrow x = 1 \text{ or } y = 1.$$

In such a case, 1 would be the rational number we are looking for. So from now on, we assume that $1 \in A_0$. Let

$$A_0 = \{1, a_1, a_2, \dots, a_n\},$$

where $a_i \neq 1$. We define

$$v_k = \frac{1+a_k}{1-a_k}.$$

We shall prove by induction on k that every element $x \in A_k$, $x \neq 1$ is representable as

$$f(v_1^{\alpha_1} v_2^{\alpha_2} \dots v_n^{\alpha_n}), \alpha_i \in \mathbb{N}_0,$$

where

$$f(x) = \frac{x-1}{x+1}.$$

Indeed:

$$f(v_k) = \frac{\frac{1+a_k}{1-a_k} - 1}{\frac{1+a_k}{1-a_k} + 1} = \frac{2a_k}{2} = a_k,$$

so the statement is true for all elements in A_0 .

Next, observe that

$$\frac{f(x) + f(y)}{1 + f(x)f(y)} = \frac{\frac{x-1}{x+1} + \frac{y-1}{y+1}}{1 + \frac{x-1}{x+1} \frac{y-1}{y+1}} = \frac{2xy - 2}{2xy + 2} = \frac{xy - 1}{xy + 1} = f(xy).$$

So for every two elements x and y in A_k with $xy \neq 1$, the induction provides representations $f(v_1^{\alpha_1} v_2^{\alpha_2} \dots v_n^{\alpha_n})$ and $f(v_1^{\beta_1} v_2^{\beta_2} \dots v_n^{\beta_n})$, so we have

$$\frac{x+y}{1+xy} = f(v_1^{\alpha_1+\beta_1} v_2^{\alpha_2+\beta_2} \dots v_n^{\alpha_n+\beta_n}).$$

That means all elements in A_{k+1} are representable in the desired form, which completes the induction. Thus

$$A_k \subseteq A \cup \{1\} \quad \text{for } k \geq 0, \quad (*)$$

where

$$A = \{f(v_1^{\alpha_1} v_2^{\alpha_2} \dots v_n^{\alpha_n}) \mid \alpha_i \in \mathbb{N}_0, \quad i = 1, 2, \dots, n\}.$$

Let p be a prime number, greater than any prime number in the prime factorization of the numerators and denominators of the rational numbers v_i , $i = 1, 2, \dots, n$. Such a prime p exists since there are finitely many v_i , and it has no representation of the form

$$p = v_1^{\alpha_1} v_2^{\alpha_2} \dots v_n^{\alpha_n},$$

since none of the factorizations of the numerators and denominators of v_i contain the prime p . Furthermore, notice that $f(x)$ is a bijection

$$f : \mathbb{R} \setminus \{-1\} \rightarrow \mathbb{R} \setminus \{1\},$$

as it is defined for every real number different from -1, and for every number $y \neq 1$ there is a unique solution $x = (1+y)/(1-y)$ to $f(x) = y$. Since p cannot be expressed as $v_1^{\alpha_1} \dots v_n^{\alpha_n}$, from the bijection we get that $f(p) \notin A$, and since $f(p) \neq 1$, it means the rational number $f(p)$ is not in any A_k , which completes the proof. \square

1.4

The lateral surface of a cylinder whose circumference is the odd prime p and whose height is a positive integer k is divided into unit squares. In each square a real number is written. For every square, the sum of the numbers written in its left and right neighbors equals the sum of the numbers written in its vertical neighbors (above and below), whenever these neighbors exist. Moreover at least one of the numbers is nonzero. Prove that $k \geq p - 1$.

Solution: Let us choose one column and consider the rows as p -dimensional vectors whose first coordinate lies in the chosen column. Let us index the vectors as a_1, a_2, \dots, a_k . We also define $a_0 = a_{k+1} = \vec{0}$ for completeness. Let C be the cyclic $p \times p$ shift matrix that has ones on the first upper diagonal with its wraparound entry and zeros everywhere else, i.e. $C_{ij} = 1$ for $(i, j) = (1, 2), (2, 3), \dots, (p-1, p), (p, 1)$. It is easy to see that C^m is the matrix with ones on the m -th upper cyclic diagonal. Hence $C^p = I$ (the identity matrix), so $C^{p-1} = C^{-1}$. The condition of the problem translates into the following relation for the vectors a_i :

$$a_{i-1} + a_{i+1} = (C + C^{-1})a_i, \quad i = 1, 2, \dots, k-1. \quad (*)$$

For the last k -th vector we have

$$a_{k-1} = (C + C^{-1})a_k,$$

and since we defined $a_{k+1} = \vec{0}$, the expression $(*)$ can be extended to $i = k$. Then from $(*)$ we have

$$\begin{aligned} a_{i+1} - Ca_i &= C^{-1}(a_i - Ca_{i-1}) \\ &= C^{-2}(a_{i-1} - Ca_{i-2}) \\ &\dots \\ &= C^{-i}(a_1 - Ca_0) = C^{-i}a_1. \end{aligned}$$

Hence by induction

$$a_{i+1} = (C^i + C^{i-2} + \dots + C^{-(i-2)} + C^{-i})a_1, \quad i = 1, 2, \dots, k.$$

Clearly $a_1 \neq \vec{0}$, otherwise the recurrence would imply $a_i = \vec{0}$ for all i , which would contradict the condition. Applying the above to $i = k$ and using $a_{k+1} = \vec{0}$, we get $Ba_1 = \vec{0}$, where

$$B = C^k + C^{k-2} + \dots + C^{-k+2} + C^{-k}.$$

Now, using $Ba_1 = \vec{0}$ we get

$$\vec{0} = CBa_1 = (CB + B)a_1 = (C^{k+1} + C^k + \dots + C^{-k+1} + C^{-k})a_1.$$

Multiplying both sides by C^k we get

$$Ta_1 = \vec{0},$$

where $T = C^0 + C^1 + \dots + C^{2k+1}$.

If we suppose that $k < p - 1$, since p is odd, we have $(2k + 2, p) = 1$, so there exist positive integers m and r such that $m(2k + 2) = rp + 1$. From $Ta_1 = \vec{0}$ and $C^p = I$ we have

$$\begin{aligned}\vec{0} &= Ta_1 = (I + C^{2k+2} + C^{2(2k+2)} + \dots + C^{(m-1)(2k+2)})Ta_1 \\ &= (T + C^{2k+2}T + \dots + C^{(m-1)(2k+2)}T)a_1 \\ &= (I + C + C^2 + \dots + C^{rp})a_1 \\ &= rpDa_1 + Ia_1,\end{aligned}$$

where D is the matrix with only 1 in all its entries. Let $a_1 = (x_1, x_2, \dots, x_p)$. Then

$$rp(x_1 + x_2 + \dots + x_p) + x_i = 0, \quad i = 1, 2, \dots, p.$$

Therefore

$$rp \sum_{j=1}^p x_j + x_i = 0 = rp \sum_{j=1}^p x_j + x_1$$

for all i , which means $x_i = x_1$ for all i . Now we have

$$\begin{aligned}0 &= rp \sum_{j=1}^p x_j + x_1 = rp \sum_{j=1}^p x_1 + x_1 \\ &= rp^2 x_1 + x_1 = (rp^2 + 1)x_1.\end{aligned}$$

Therefore $x_1 = 0$ and so $x_i = x_1 = 0$ for $i = 1, 2, \dots, p$, implying that $a_1 = \vec{0}$, which is a contradiction. Thus $k \geq p - 1$ and the problem is solved \square

1.5

A sequence of polynomials is defined in the following way:

$$p_0(x) = x,$$

$$p_n(x) = 1 + p_0(x)p_1(x) \cdots p_{n-1}(x), \quad n = 1, 2, \dots$$

Prove that for $n > 0$ the number of odd coefficients of $p_n(x)$ is one more than some integer power of the number 2.

Solution: We have

$$p_1(x) = 1 + x$$

and

$$p_n(x) = 1 + (p_{n-1}(x) - 1)p_{n-1}(x) = p_{n-1}(x)^2 - p_{n-1}(x) + 1.$$

From now on, we work modulo 2. First, notice that for every polynomial $q(x)$ we have

$$\begin{aligned} q(x)^2 &= (a_0 + a_1x + \cdots + a_nx^n)^2 \\ &= a_0^2 + a_1^2x^2 + \cdots + a_n^2x^{2n} + 2 \sum_{i \neq j} a_i a_j x^{i+j} \\ &= a_0 + a_1x^2 + \cdots + a_nx^{2n} = q(x^2). \end{aligned}$$

Therefore, from the recursive relation for p_n , we have

$$p_n(x) = p_{n-1}(x)^2 - p_{n-1}(x) + 1 = p_{n-1}(x^2) + p_{n-1}(x) + 1. \quad (*)$$

By induction, it is seen that the degree of $p_n(x)$ is exactly 2^{n-1} , that it always has a constant term 1, and that besides x^0 it has nonzero coefficients only in front of x^{2^i} for some $i \geq 0$. Let the coefficient in front of x^{2^i} be $a_i^{(n)}$ for $i = 0, 1, \dots, n-1$, i.e.

$$p_n(x) = 1 + a_0^{(n)}x^{2^0} + a_1^{(n)}x^{2^1} + a_2^{(n)}x^{2^2} + \cdots + a_{n-1}^{(n)}x^{2^{n-1}}.$$

From (*), we know that:

- the coefficient in front of $x = x^{2^0}$ in $p_n(x)$ is only contributed by $p_{n-1}(x)$, so $a_0^{(n)} = a_0^{(n-1)}$;
- the coefficient in front of $x^{2^{n-1}}$ in $p_n(x)$ is only contributed by $p_{n-1}(x^2)$, so $a_{n-1}^{(n)} = a_{n-2}^{(n-1)}$;
- for $1 \leq i \leq n-2$, the coefficient in front of x^{2^i} in $p_n(x)$ is the sum of $a_{i-1}^{(n-1)}$, owing to $p_{n-1}(x^2)$, and $a_i^{(n-1)}$, owing to $p_{n-1}(x)$, so $a_i^{(n)} = a_{i-1}^{(n-1)} + a_i^{(n-1)}$.

But that is the exact recurrence relation that the binomial coefficients satisfy, namely

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Therefore, by induction, we get

$$a_i^{(n)} = \binom{n-1}{i}.$$

Thus the number of odd coefficients of $p_n(x)$ is 1 plus the number of odd binomial coefficients of order $n-1$. But from Lucas' theorem, $\binom{n-1}{i}$ is odd exactly when every digit in the binary representation of $n-1$ is at least the corresponding digit in the binary representation of i . That means $a_i^{(n)} \neq 0$ for precisely those i that have 0 wherever $n-1$ has 0 and 0 or 1 whenever $n-1$ has 1. The number of such i is thus

$$2^{S(n-1)},$$

where $S(n-1)$ is the number of 1s in the binary representation of $n-1$. That completes the proof. \square

1.6

Prove that the following holds for every positive integer n :

$$\left| \left\{ \frac{n}{1} \right\} - \left\{ \frac{n}{2} \right\} + \left\{ \frac{n}{3} \right\} - \cdots + (-1)^{n-1} \left\{ \frac{n}{n} \right\} \right| < \sqrt{2n}.$$

Solution: For $n = 1, 2, 3, 4$ we verify directly, so we can assume $n > 4$. Let us take an integer $k \in [\sqrt{n}, n]$. We know that $k > 2$. We shall split the sum in two parts and bound each part separately:

$$A_k = \left\{ \frac{n}{1} \right\} - \left\{ \frac{n}{2} \right\} + \left\{ \frac{n}{3} \right\} - \cdots + (-1)^k \left\{ \frac{n}{k-1} \right\}$$

and

$$B_k = \left\{ \frac{n}{k} \right\} - \left\{ \frac{n}{k+1} \right\} + \left\{ \frac{n}{k+2} \right\} - \cdots + (-1)^{n-k} \left\{ \frac{n}{n} \right\} = C_k - D_k,$$

where

$$C_k = \left(\frac{n}{k} - \frac{n}{k+1} + \cdots + (-1)^{n-k} \frac{n}{n} \right),$$

$$D_k = \left(\left\lfloor \frac{n}{k} \right\rfloor - \left\lfloor \frac{n}{k+1} \right\rfloor + \cdots + (-1)^{n-k} \left\lfloor \frac{n}{n} \right\rfloor \right).$$

With respect to C_k we have:

$$C_k = \frac{n}{k} - \left(\frac{n}{k+1} - \frac{n}{k+2} \right) - \cdots \leq \frac{n}{k},$$

since all the terms in the brackets are positive (no matter whether the last bracket contains one or two terms). Furthermore, equality holds only when $k = n$, when the bracketed terms are missing.

Furthermore,

$$C_k = \left(\frac{n}{k} - \frac{n}{k+1} \right) + \left(\frac{n}{k+2} - \frac{n}{k+3} \right) + \cdots > 0,$$

so we conclude $0 < C_k \leq \frac{n}{k}$.

For D_k we have

$$D_k = \left(\left\lfloor \frac{n}{k} \right\rfloor - \left\lfloor \frac{n}{k+1} \right\rfloor \right) + \left(\left\lfloor \frac{n}{k+2} \right\rfloor - \left\lfloor \frac{n}{k+3} \right\rfloor \right) + \cdots$$

For $k \geq \sqrt{n}$ and for $i = 0, 1, \dots$ we have

$$\frac{n}{k+2i} - \frac{n}{k+2i+1} < \frac{n}{k(k+1)} < 1,$$

and so

$$\left\lfloor \frac{n}{k+2i} \right\rfloor - \left\lfloor \frac{n}{k+2i+1} \right\rfloor \in \{0, 1\}.$$

Notice that among the inequalities

$$\left\lfloor \frac{n}{k+i} \right\rfloor - \left\lfloor \frac{n}{k+i+1} \right\rfloor \leq 1$$

exactly $\left\lfloor \frac{n}{k} \right\rfloor - 1$ are strict, since the telescopic sum of all those terms for $i = 0, 1, \dots, n-k-1$ is exactly

$$\left\lfloor \frac{n}{k+0} \right\rfloor - \left\lfloor \frac{n}{n} \right\rfloor = \left\lfloor \frac{n}{k} \right\rfloor - 1.$$

Then at most $\left\lfloor \frac{n}{k} \right\rfloor - 1$ of the inequalities

$$\left\lfloor \frac{n}{k+2i} \right\rfloor - \left\lfloor \frac{n}{k+2i+1} \right\rfloor \leq 1$$

are strict too (the even indices being a subset of all), and therefore

$$0 \leq D_k \leq \left\lfloor \frac{n}{k} \right\rfloor - 1 < \frac{n}{k},$$

where $D_k = 0$ is not possible for $k = n$, because then $D_k = 1$. Finally

$$|B_k| = |C_k - D_k| < \frac{n}{k}$$

and therefore the inequality is strict.

For A_k we have

$$-\left\{ \frac{n}{2} \right\} - \left\{ \frac{n}{4} \right\} - \dots \leq A_k \leq \left\{ \frac{n}{3} \right\} + \left\{ \frac{n}{5} \right\} + \dots$$

From

$$0 \leq \left\{ \frac{n}{i} \right\} \leq \frac{i-1}{i} \leq \frac{k-2}{k-1}$$

it follows

$$|A_k| \leq \left\lceil \frac{k-2}{2} \right\rceil \frac{k-2}{k-1} \leq \left(\frac{k-2}{2} + \frac{1}{2} \right) \frac{k-2}{k-1} = \frac{k-2}{2}.$$

Since the sum in the statement is equal to $A_k + (-1)^{k+1}B_k$ and

$$|A_k \pm B_k| \leq |A_k| + |B_k| < \frac{k-2}{2} + \frac{n}{k},$$

to complete the proof we need to find some $k \in [\sqrt{n}, n]$ such that:

$$\frac{k-2}{2} + \frac{n}{k} \leq \sqrt{2n}.$$

That last inequality is equivalent to

$$k^2 - (2 + \sqrt{8n})k + 2n \leq 0,$$

and so k must lie between the roots of the quadratic equation, i.e. it must satisfy

$$k \in [\gamma - \delta, \gamma + \delta],$$

where $\gamma = \sqrt{2n} + 1$ and $\delta = \sqrt{\sqrt{8n} + 1}$.

We shall show that the choice of $k = \lceil \sqrt{2n} \rceil$ satisfies both requirements. First, $k < 1 + \sqrt{2n} = \gamma < \gamma + \delta$. Next, $k \geq \sqrt{2n} = \gamma - 1 > \gamma - \delta$, since $\delta > 1$. That gives us $k \in [\gamma - \delta, \gamma + \delta]$. That $k > \sqrt{n}$ follows directly from $k \geq \sqrt{2n} > \sqrt{n}$. And finally, since $k < \gamma$, to prove $k \leq n$, it suffices to show that $\gamma < n$. But

$$\gamma < n \leftrightarrow 2n < (n - 1)^2 \leftrightarrow 0 < n(n - 4) + 1,$$

which is true for $n \geq 4$. Therefore $k \in [\sqrt{n}, n]$ and all requirements are satisfied. \square

1.7

Prove that

- a) there does not exist a function $f : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}$,
 b) there exists a function $f : \mathbb{Q}^+ \times \mathbb{Q}^+ \rightarrow \mathbb{Q}$,
 such that for all $a < b < c$ in the domain of f holds:

$$f(a, c) < f(a, b) < f(b, c).$$

Proof a) Suppose that such a function exists. For every positive real number x , using $a = x, b = 2x, c = 3x$, we get

$$f(x, 3x) < f(x, 2x).$$

For every x , consider the interval $\Delta(x) = (f(x, 3x), f(x, 2x))$. We shall see that these intervals are pairwise disjoint. Let us take any $x < y$ and examine the three possibilities:

- Case 1. $y > 2x$. We have $x < 2x < y < 3y$. Taking $a = x, b = 2x, c = 3y$, we obtain $f(x, 2x) < f(2x, 3y)$; and taking $a = 2x, b = y, c = 3y$, we obtain $f(2x, 3y) < f(y, 3y)$. Combining the two inequalities gives $f(x, 2x) < f(2x, 3y) < f(y, 3y)$.
- Case 2. $y = 2x$. As in the previous case, we take $a = x, b = 2x, c = 3y$ to obtain $f(x, 2x) < f(2x, 3y)$, but this time $f(2x, 3y) = f(y, 3y)$, so now $f(x, 2x) < f(2x, 3y) = f(y, 3y)$.
- Case 3. $y < 2x$. We have $x < y < 2x < 3y$. Taking $a = x, b = y, c = 2x$, we obtain $f(x, 2x) < f(x, y)$; and taking $a = x, b = y, c = 3y$, we obtain $f(x, y) < f(y, 3y)$. Combining the two inequalities gives $f(x, 2x) < f(x, y) < f(y, 3y)$.

In all examined cases, we have $f(x, 2x) < f(y, 3y)$, and so the intervals $\Delta(x)$ and $\Delta(y)$ do not intersect. But each interval $\Delta(x)$ contains infinitely many rational numbers. Choose a rational number from each interval and denote it by $q(x)$. Since the intervals are pairwise disjoint, all $q(x)$ are different, which means q is an injection from $\mathbb{R}^+ \rightarrow \mathbb{Q}$. But, as is well known, this is impossible. Consequently, a function with the sought properties does not exist.

Proof b) Notice that it does not matter how we define $f(x, y)$ for $x \geq y$. Our goal is to construct a function f such that: (a) $f(a, x)$ is decreasing in x for every fixed a and all $x > a$; and (b), the image $F_a = \{f(a, x) : x > a\}$ is to the left of the image $F_b = \{f(b, x) : x > b\}$ for every $a < b$. The second condition is the hard one, and it requires us to do in \mathbb{Q} what we could not do in \mathbb{R} earlier, namely to construct a set of non-overlapping intervals where between every two intervals from the set there is a third one from the same set.

Part 1: Constructing the intervals. Consider all sequences p_1, p_2, \dots, p_{n-1} where $p_i \in \{0, 2\}$, and for each such sequence define l and r as follows:

$$l = (0.p_1p_2 \cdots p_{n-1}1)_3; \quad r = (0.p_1p_2 \cdots p_{n-1}2)_3,$$

where the subscript 3 denotes *ternary representation*. In other words,

$$l = \sum_{i=1}^{n-1} \frac{p_i}{3^i} + \frac{1}{3^n}, \quad r = \sum_{i=1}^{n-1} \frac{p_i}{3^i} + \frac{2}{3^n}.$$

Let C be the set of all intervals (l, r) for l and r defined in the above fashion. We shall show that between every two intervals in C there exists a third interval from C ; therefore the intervals in C must be disjoint.

Let us take two different intervals (l_1, r_1) and (l_2, r_2) . Clearly $r_1 \neq r_2$, else we would have $l_1 = l_2$ and the intervals would be the same. Without loss of generality, assume $r_1 < r_2$. Then their ternary representations must differ in some digit, where the digit of r_2 is larger, and this digit cannot be the last one (as it is always 2). If r_2 is written with n symbols, then

$$r_2 - r_1 > \frac{2}{3^{n-1}} - \left(\frac{2}{3^n} + \frac{2}{3^{n+1}} + \cdots \right) = \frac{1}{3^{n-1}}.$$

Let

$$r_3 = r_2 - \frac{2}{3^n} + \frac{2}{3^{n+1}}, \quad l_3 = r_2 - \frac{2}{3^n} + \frac{1}{3^{n+1}}.$$

Clearly (l_3, r_3) is from C , as the ternary representations of l_3 and r_3 are identical up to the last digit, which is 1 for l_3 and 2 for r_3 . Therefore,

$$r_2 > l_2 = r_2 - \frac{1}{3^n} > r_3 > l_3 > r_2 - \frac{1}{3^{n-1}} > r_1 > l_1,$$

which shows that (l_3, r_3) sits between (l_1, r_1) and (l_2, r_2) .

Furthermore, for every interval I in C , we could choose a sufficiently large n so that the following two intervals (also in C) sit to the left and right of I :

$$\left(\frac{1}{3^n}, \frac{2}{3^n} \right) \text{ and } \left(1 - \frac{2}{3^n}, 1 - \frac{1}{3^n} \right).$$

Now we define a map $\Delta : \mathbb{Q}^+ \rightarrow C$. Let us list the positive rational numbers as a sequence: q_1, q_2, \dots and define their corresponding intervals $\Delta(q_i)$ sequentially. First, we assign to q_1 an arbitrary interval $\Delta(q_1)$ from C . Thereafter, for every $m > 1$, we assign $\Delta(q_m) \in C$ as follows:

- if q_m is larger than all previously listed numbers, we choose an interval from C that is to the right of all intervals $\Delta(q_i)$ and assign it to q_m .
- if q_m is smaller than all previously listed numbers, we choose an interval from C that is to the left of all intervals $\Delta(q_i)$ and assign it to q_m .
- if q_m is neither the smallest nor the largest among the previously listed numbers, let $q_i < q_m < q_j$ be the two numbers among them immediately adjacent to q_m . In that case, we pick an interval from C that lies between $\Delta(q_i)$ and $\Delta(q_j)$ and assign it to q_m .

Part 2: Construction of f . Now that our map Δ is defined, we construct f as follows. For every $a \in \mathbb{Q}^+$ we take our $\Delta(a) = (l_a, r_a)$ and define $f(a, x)$ as follows:

$$f(a, x) = \begin{cases} 1983 & \text{if } x \leq a \\ l_a + \frac{r_a - l_a}{x+1} & \text{if } x > a \end{cases}$$

Thus the function f is well defined for all $(a, x) \in \mathbb{Q}^+ \times \mathbb{Q}^+$ and its values lie in \mathbb{Q} , since l_a, r_a, x are positive rational numbers. We shall show that it satisfies the condition.

Let $a < b < c$ be any positive rational numbers. Since $f(a, x)$ is a decreasing function on $x > a$, we have $f(a, c) < f(a, b)$, so the first inequality is satisfied.

Next, notice that for all $x > a$, $l_a < f(a, x) < r_a$, so $f(a, x) \in \Delta(a)$, in particular $f(a, b) \in \Delta(a)$. Likewise, $f(b, c) \in \Delta(b)$. But since $a < b$, from the construction of the map Δ , we know that $\Delta(a)$ sits entirely to the left of $\Delta(b)$. Therefore $f(a, b) < f(b, c)$. The second inequality is also satisfied, and our construction is complete. \square

2 Combinatorics

2.1

Let A, B be lattice points with positive coordinates such that $OA = OB$. Prove that the circles with diameters OA and OB contain the same number of lattice points with positive coordinates.

Solution: Let $A = (m, n)$ and $B = (a, b)$ and the two circles be k_1 and k_2 respectively. The condition $(x, y) \in k_1$ is equivalent to

$$\begin{aligned}x^2 + y^2 + (m - x)^2 + (n - y)^2 &= m^2 + n^2 \\ \iff (2x - m)^2 + (2y - n)^2 &= m^2 + n^2.\end{aligned}$$

Since

$$a^2 + b^2 = OB^2 = OA^2 = m^2 + n^2,$$

either (a) $m \equiv a, n \equiv b \pmod{2}$ or (b) $m \equiv b, n \equiv a \pmod{2}$. Indeed, if OA^2 is odd, then (a, b) and (m, n) are pairs of numbers of different parity, and if OA^2 is even, then a, b, m, n are of the same parity.

In the case (a) there is a bijection between the lattice points on the two circles:

$$f : \{(x, y) \mid (x, y) \in k_1\} \rightarrow \{(x', y') \mid (x', y') \in k_2\},$$

given by

$$f(x, y) = \left(x + \frac{a - m}{2}, y + \frac{b - n}{2} \right).$$

Similarly, in case (b), we can define the bijection as follows:

$$f(x, y) = \left(y + \frac{a - n}{2}, x + \frac{b - m}{2} \right).$$

Thus we proved that k_1 and k_2 have the same number N of integer points.

Let us take the positive integer points (x, y) , $x, y > 0$, of k_1 . We have

$$x^2 + y^2 = mx + ny.$$

If $x > m$, then

$$0 < (x - m)x = y(n - y) \Rightarrow y < n \quad (\text{type 1}).$$

If $x < m$, then

$$y(y - n) = x(m - x) > 0 \Rightarrow y > n \quad (\text{type 2}).$$

If $x = m$, then

$$y = n \quad (\text{type 3}).$$

Clearly there are no points on k_1 with both negative coordinates, owing to $x^2 + y^2 = mx + ny$. The number of points $(u, v) \in k_1$ with positive u and negative v equals the number

of positive points of type 2, since there is a bijection between the two sets given by $(u, v) \leftrightarrow (x, n - v)$. Likewise, the points $(u, v) \in k_1$ with positive v and negative u correspond to the positive points of type 1, as demonstrated by the bijection $(u, v) \leftrightarrow (m - x, y)$.

Including the points $(u, v) \in k_1$ with $uv = 0$, namely $(0, 0)$, $(m, 0)$, $(0, n)$, and the one point of type 3, we have:

$$N = 2(\# \text{type 1}) + 2(\# \text{type 2}) + 3 + 1.$$

Hence the number of positive points is

$$(\# \text{type 1}) + (\# \text{type 2}) + 1 = \frac{N - 4}{2} + 1 = \frac{N}{2} - 1.$$

The same holds for the second circle, which completes the proof.

Geometrically the correspondence between the circles is a translation in case (a) and a reflection across the line $x = y$ followed by a translation in case (b). \square

2.2

Given is a sequence a_1, a_2, \dots, a_n of distinct real numbers, such that there does not exist a decreasing subsequence of length $k + 1$. Prove that the number of pairs $\{i, j\}$ for which $i < j$ and $a_i > a_j$ does not exceed

$$\frac{k-1}{2k}n^2.$$

Solution: A pair (i, j) with $i < j$ and $a_i > a_j$ is called an *inversion*. For every $i = 1, 2, \dots, n$, define $r(i)$ as the maximal possible number r for which there exists a sequence

$$i_1 < i_2 < \dots < i_{r-1} < i, \quad a_{i_1} > a_{i_2} > \dots > a_{i_{r-1}} > a_i.$$

From the condition we have $r(i) \leq k$. Let us take an arbitrary inversion (i, j) and let $r = r(j)$. We have

$$i_1 < i_2 < \dots < i_{r-1} < j, \quad a_{i_1} > a_{i_2} > \dots > a_{i_{r-1}} > a_j,$$

which implies that $r(i) < r(j)$. Let us divide the indices into k sets $A(1), A(2), \dots, A(k)$, where $i \in A(r(i))$. We have

$$|A(1)| + |A(2)| + \dots + |A(k)| = n,$$

and for every two elements i and j from one and the same set $A(r)$ the pair (i, j) is not an inversion. Therefore the number of all pairs which are not inversions is at least

$$\binom{|A(1)|}{2} + \binom{|A(2)|}{2} + \dots + \binom{|A(k)|}{2}.$$

Hence the number of inversions does not exceed

$$\binom{n}{2} - \left(\binom{|A(1)|}{2} + \binom{|A(2)|}{2} + \dots + \binom{|A(k)|}{2} \right).$$

Finally, from the inequality of Cauchy, we obtain:

$$\begin{aligned} \binom{n}{2} - \sum_{r=1}^k \binom{|A(r)|}{2} &\leq \frac{n(n-1) - \sum_{r=1}^k |A(r)|^2 + \sum_{r=1}^k |A(r)|}{2} \\ &\leq \frac{n^2 - \frac{n^2}{k}}{2} = \frac{k-1}{2k}n^2. \end{aligned}$$

2.3

There is a 6 by 21 table (a_{ij}) with a total of 18 zeros, 18 ones, \dots , 18 sixes. For any $k, l \in \{1, 2, \dots, 6\}$ and $p, q \in \{1, 2, \dots, 21\}$ we have

$$a_{k,p} - a_{l,p} \equiv a_{k,q} - a_{l,q} \pmod{7}.$$

Prove that every row contains 3 zeros, 3 ones, \dots , 3 sixes.

Solution: Let us construct a new table (b_{ij}) , where

$$b_{ij} = \omega^{a_{ij}}, \quad \omega = \cos \frac{2\pi}{7} + i \sin \frac{2\pi}{7}.$$

Let the sum of the numbers in the k -th row be s_k , $k = 1, 2, \dots, 6$. From the condition we have

$$a_{k,p} - a_{1,p} \equiv a_{k,1} - a_{1,1} \equiv m_k \pmod{7}$$

for some integer m_k with $0 \leq m_k \leq 6$ and all $p = 1, 2, \dots, 21$, and hence

$$s_k = s_1 \omega^{m_k}.$$

Let

$$q(x) = 1 + x + \dots + x^6.$$

Since each of the numbers $0, 1, \dots, 6$ appears 18 times in the entire table, summing over all rows we get

$$s_1(\omega^0 + \omega^1 + \dots + \omega^6) = \sum_{k=1}^6 s_k = 18q(\omega) = 0.$$

Let

$$p(x) = x^{m_1} + x^{m_2} + \dots + x^{m_6}$$

and

$$s(x) = \sum_{i=1}^{21} x^{a_{1i}},$$

so

$$s(\omega) = s_1.$$

Thus

$$s(\omega)p(\omega) = 0,$$

i.e. ω is a root of either $s(x)$ or $p(x)$.

Now we use the well-known fact that the polynomial $q(x)$ is the 7th cyclotomic polynomial and is irreducible over \mathbb{Q} .

If we assume that $p(\omega) = 0 = q(\omega)$, then from the irreducibility of $q(x)$ it follows that $q(x)$ divides $p(x)$. From $\deg p(x) \leq 6 = \deg q(x)$ it follows that $p(x) = kq(x)$ for some integer k . But then

$$7k = q(1)k = p(1) = 6,$$

which is impossible. Consequently $p(\omega) \neq 0$ and so it must be the case that $s(\omega) = 0$.

Since $s(\omega) = 0$ and $q(x)$ is the minimal polynomial of ω over \mathbb{Q} , it follows that $q(x)|s(x)$, and since $\deg s \leq 6 = \deg q$, we must have

$$s(x) = kq(x)$$

for some integer k . Evaluating at $x = 1$, we obtain

$$21 = s(1) = kq(1) = 7k,$$

i.e. $k = 3$ and therefore

$$s(x) = 3 + 3x + 3x^2 + 3x^3 + 3x^4 + 3x^5 + 3x^6.$$

This proves that in the first row of the table there are 3 zeros, 3 ones, \dots , 3 sixes. The same holds for the remaining rows, which completes the proof. \square

2.4

We are given n switches and n lamps. We know that each switch connects to exactly one lamp, but not which switch connects to which lamp. At each move, we are allowed to flip simultaneously any switches and observe which lamps change state. Find the minimum number $f(n)$ of moves that guarantees we can switch all lamps OFF regardless of their original ON/OFF state.

Solution: The general idea is to show that after each move, there is always the possibility that at least half of the work remains undone, with slightly more ON lamps than OFF lamps remaining until the end, when we will need one extra move to turn off the last lamp.

Let $t_c = \lceil \log_2 n \rceil$. Suppose that at the start exactly $\lceil n/2 \rceil$ (half or slightly more when n is odd) of the lamps are ON. We shall prove the following lemma:

Lemma. After any $0 \leq t \leq t_c$ moves, regardless of how we play, it may still be possible that there exist sets L_t of lamps and S_t of switches such that:

1. $|S_t| \geq |L_t| = n_t \geq \lceil n/2^t \rceil \geq 1$;
2. on every move, either all switches in S_t were flipped together or none were flipped; and
3. $k_t = \lceil n_t/2 \rceil \geq 1$ of the n_t lamps in L_t are ON.

Proof: For $t = 0$, all requirements are true with S_0 and L_0 consisting of all switches and all lamps respectively. By induction on t , consider $1 \leq t \leq t_c$. We have $|L_{t-1}| = n_{t-1} \geq 2$. Suppose after our t -th move we observe exactly m of the lamps in L_{t-1} change state. Consider two cases:

Case 1. $m < k_{t-1}$, so less than half of the lamps in L_{t-1} changed state. Let $L_t \subseteq L_{t-1}$ be the set of $n_t = n_{t-1} - m$ lamps that *did not change state*, and let $S_t \subseteq S_{t-1}$ be the switches we did not flip. Each lamp in L_t must necessarily connect to a switch in S_t , so $|S_t| \geq |L_t| = n_t$. Since $n_t \geq n_{t-1}/2 \geq n/2^t$, requirement 1 is satisfied. Furthermore, none of the switches in S_t were flipped at the t -th move, and since they are a subset of S_{t-1} , Requirement 2 is satisfied as well.

Finally, to satisfy the third requirement, notice that since all switches in S_{t-1} were always flipped together or not flipped on every move, all lamps in L_{t-1} have exhibited exactly the same history on the first $t - 1$ moves, because their controlling switches all lie in S_{t-1} and were treated identically on every previous move. From the observations so far, any matching of those lamps to distinct switches in S_{t-1} remains consistent with what we have seen. So L_t could consist of any n_t lamps from L_{t-1} . As a result, we may get exactly k_t lamps from among the k_{t-1} in L_{t-1} that were ON (and still are) and $n_t - k_t$ from the set of $n_{t-1} - k_{t-1}$ lamps that were OFF (and still are). Since $n_{t-1} \geq n_t$, this is feasible since $k_t \leq \lceil n_{t-1}/2 \rceil = k_{t-1}$ and $n_t - k_t = \lfloor n_t/2 \rfloor \leq \lfloor n_{t-1}/2 \rfloor = n_{t-1} - k_{t-1}$.

Case 2. $m \geq k_{t-1}$, so at least half of the lamps in L_{t-1} changed state. Let L_t be a set of $n_t = k_{t-1} < n_{t-1}$ chosen from the lamps of L_{t-1} that *did change state*. That $k_{t-1} < n_{t-1}$ follows from $t \leq t_c$ and $n_{t-1} \geq 2$. Let $S_t \subseteq S_{t-1}$ be the switches we flipped. Since all

switches in S_t were flipped on the t -th move and since $S_t \subseteq S_{t-1}$, Requirement 2 holds. Since $n_t = k_{t-1} = \lceil n_{t-1}/2 \rceil \geq \lceil n/2^t \rceil$, Requirement 1 holds too.

Again, to satisfy the third requirement, notice that under the same argument as before, L_t could comprise any n_t lamps from L_{t-1} . As a result we may get exactly k_t from among the $n_{t-1} - k_{t-1}$ lamps that were OFF (and are now ON) and $n_t - k_t$ lamps from the k_{t-1} that were ON and are now OFF. From the strict inequality $n_{t-1} > k_{t-1} = n_t$, this is feasible since $k_t = \lceil n_t/2 \rceil = \lfloor (n_t+1)/2 \rfloor \leq \lfloor n_{t-1}/2 \rfloor = n_{t-1} - k_{t-1}$ and $n_t - k_t = \lfloor n_t/2 \rfloor \leq \lfloor n_{t-1}/2 \rfloor \leq k_{t-1}$.

That shows that in both cases we can continue the induction, and the lemma is proved.

Using $t = t_c$ we see that there will still exist a set L_{t_c} containing at least one lamp that is still ON, and so we need at least one more move to turn it off, hence $f(n) \geq 1 + t_c$.

To prove the reverse inequality, namely that $1 + t_c$ moves always suffice, let's number the switches from 0 to $n - 1$ and consider their binary representation, which uses at most t_c digits. At every move $t = 1, 2, \dots, t_c$, we flip the switches that have 1 in their t -th digit. Each lamp will flip exactly in the pattern corresponding to the binary code of its switch, so after t_c moves we know which switch controls which lamp. We may need one last move to switch off any lamps that remain ON. So finally, $f(n) \leq 1 + t_c$ □

2.5

Let there be several distinct words of p ones and q zeros ($p > q$). Prove that we can pick a 1 from each word and replace it with 0 so that the obtained words of $p - 1$ ones and $q + 1$ zeros are again distinct.

Solution: We apply induction on $p + q$. If $p + q = 1$, then $p = 1$, $q = 0$. There is only one such word, 1, and replacing the one with zero results in a single word, so the statement is correct. Let us now assume that $p + q > 1$ and that we have proved the statement for all shorter words. Consider two cases:

Case 1. $p > q + 1$. Divide the given words into two groups: those beginning with 1 and those beginning with 0.

Consider the first group and remove the first symbol from each word. The resulting shortened words contain $p - 1$ ones and q zeros. Because $p > q + 1$, we have $p - 1 > q$, so the induction hypothesis applies to these shortened words: we can replace a one in each of them so that they remain distinct. Restoring the initial symbol 1 to each shortened word yields words that are still distinct.

The same argument applies to the second group. After removing the first symbol, the shortened words contain p ones and $q - 1$ zeros, and since $p > q - 1$, the induction hypothesis again applies. Restoring the initial symbol 0 yields distinct words within this group as well.

Finally, every word obtained from the first group begins with 1, while every word obtained from the second group begins with 0. Hence the resulting words from the two groups are distinct from each other.

Case 2. $p = q + 1$. In this case, the given words are a subset of the set A of all words with $q + 1$ ones and q zeros. We shall prove the statement for the entire set A .

Let B be the set of all words with q ones and $q + 1$ zeros. Changing any of the $q + 1$ ones to zero for any given word in A results in a word from B , so each word $x \in A$ maps to a set $\Gamma(x) \subseteq B$ of size $q + 1$. Vice versa, switching any of the $q + 1$ zeros in B to one for any word $y \in B$ results in a distinct word $x \in A$ with $y \in \Gamma(x)$. Therefore, each word in B belongs to exactly $q + 1$ of the sets $\Gamma(x)$. Let us take an arbitrary k -element subset $X = \{x_1, x_2, \dots, x_k\}$ of A and consider the union

$$\Gamma(X) = \Gamma(x_1) \cup \Gamma(x_2) \cup \dots \cup \Gamma(x_k).$$

Let the number of pairs $\{x, y\}$, $x \in X$, $y \in \Gamma(X)$ be $u(X)$. On one hand,

$$u(X) = |X|(q + 1) = k(q + 1).$$

On the other hand, every element of $\Gamma(X)$ participates in at most $q + 1$ such pairs, whence $|\Gamma(X)|(q + 1) \geq u(X) = k(q + 1)$, and therefore $|\Gamma(X)| \geq k$.

By Hall's theorem, the family $\{\Gamma(x) : x \in A\}$ has a system of distinct representatives. Thus there exists an injective map $f : A \rightarrow B$ such that $f(x) \in \Gamma(x)$ for every $x \in A$. This completes the proof. \square

2.6

For every integer x, a, b , we shall say that $x \in ((a, b))$ if $x \neq b$ and

$$(x - a)(x - b)(a - b) \geq 0.$$

A “nice” partition of $\{1, 2, \dots, 2n\}$ into pairs $(a_1, b_1), (a_2, b_2), \dots, (a_n, b_n)$ shall be one for which

- (i) $\exists p, 1 \leq p \leq 2n : p \in ((a_i, b_i))$ for $i = 1, 2, \dots, n$;
- (ii) $\exists q, 1 \leq q \leq 2n : q \notin ((a_i, b_i))$ for $i = 1, 2, \dots, n$.

Prove that the number of nice partitions is $2n \times n!$.

Solution: Let us arrange the numbers $1, 2, \dots, 2n$ on a circle clockwise. From here onward, we consider the indices modulo $2n$, i.e. by point $2n + 1$ we will mean point 1, point $2n + 2$ will be point 2, point 0 will be $2n$ etc.

We draw an arrow from a_i to b_i for each i and call it arrow i . We define the “right” side of arrow i as the set $\{b_i, b_i + 1, \dots, a_i - 1\}$, i.e. all the numbers that lie on the clockwise arc from b_i to a_i , including b_i but excluding a_i . Analogously, we define “left” side for every arrow as the complementary set $\{a_i, a_i + 1, \dots, b_i - 1\}$.

We shall make use of the following lemma:

Lemma. $x \in ((a, b))$ is equivalent to x being to the left of the arrow from a to b .

Proof: Examine two cases: $a < b$ and $b < a$.

- Case 1. $1 \leq a < b \leq 2n$. The points to the left of the arrow from a to b are $a, a + 1, \dots, b - 1$. For each of those points x , either $x = a$ or x is strictly in between a and b , so $(x - a)(x - b) \leq 0$. With $a - b < 0$, that gives us $x \in ((a, b))$. Conversely, if $x \in ((a, b))$, then $x \neq b$ and $(x - a)(x - b) \leq 0$, so either $x = a$ or x is between a and b . But that is precisely the set of points to the left of the arrow.
- Case 2. $1 \leq b < a \leq 2n$. The points to the left of the arrow from a to b are $a, a + 1, 2n, 1, 2, \dots, b - 1$. For each of those points x , either $x < b$ or $x \geq a$. That means that $(x - a)(x - b) \geq 0$, and with $a - b > 0$, we get $x \in ((a, b))$. Conversely, if $x \in ((a, b))$, then $x \neq b$ and $(x - a)(x - b) \geq 0$, so either $x = a$ or $x > a > b$ or $x < b < a$, and all those points are to the left of the arrow.

That completes the proof of the lemma. Notice that (i) and (ii) in the condition are equivalent to $f(p) = n$ and $f(q) = 0$.

For each point x on the circle, define $f(x)$ as the number of arrows that have x to their left.

Now let us consider any two neighboring numbers $a = x$, and $b = x + 1$.

If there is an arrow connecting a and b , then for every other arrow they are simultaneously to the left or simultaneously to the right. Now, with respect to the arrow connecting them, exactly one of a and b (the one where the arrow originates) is to the left of it, and therefore

$$f(a) - f(b) = \pm 1.$$

Likewise, if there is no arrow connecting a and b , then a and b are simultaneously to the left or right of any arrow not incident with either a or b . Let us see that the same holds for the arrow incident with a . If it originates from a , then both a and b are to its left; and if it ends in a , then neither a nor b are to its left. In either case, that arrow contributes equally to $f(a)$ and $f(b)$.

The only difference between $f(a)$ and $f(b)$ comes from the arrow incident with b . If it originates in b , then a is to its right and b is to its left, so it contributes $+1$ to $f(b)$ only. And if it ends in b , then a is to its left and b to its right, so it contributes to $+1$ to $f(a)$ only. That means that

$$f(a) - f(b) = \pm 1,$$

where the sign is determined by whether the arrow incident with b originates or ends in b .

Let $\Delta(x) = f(x+1) - f(x) = \pm 1$. From the above observations, we see that if the path from one number a to another number b along the circle (in either direction) is of length m , then $|f(a) - f(b)| \leq m$ since it is a sum of m deltas, each equal to ± 1 .

Thus from $f(p) = n$ and $f(q) = 0$ it follows that the path from p to q cannot be shorter than n , so they must be opposite points on the circle. Furthermore, since

$$n = f(p) - f(q) = \Delta(q) + \Delta(q+1) + \cdots + \Delta(p-1),$$

and all those n deltas are ± 1 , then they must necessarily be all $+1$. Therefore

$$f(q+i) = f(q) + \Delta(q) + \Delta(q+1) + \cdots + \Delta(q+i-1) = i$$

for $i = 1, 2, \dots, n-1$. Similarly, we can see that $f(p+i) = n-i$ for $i = 1, 2, \dots, n-1$. That means that for $x \neq p, x \neq q$ we have $0 < f(x) < n$. One consequence is that p is the only point x for which $f(x) = n$ and q is the only point x for which $f(x) = 0$.

Furthermore, since $f(p) = n$ and $f(q) = 0$, it means that for each of the n arrows, p must be to its left and q to its right. That means all arrows must originate in

$$P = \{p+1, p+2, \dots, q\}$$

and end in

$$Q = \{q+1, q+2, \dots, p\}.$$

Conversely, any choice of two opposite points p and q and any bijection from P to Q produce a nice partition. To see why, observe that the arrows representing the bijection originate in P and end in Q , so they have p on their left and q on their right, which means p satisfies (i) and q satisfies (ii), implying the partition is nice.

There are $2n$ ways in which the ordered pair of opposite points (p, q) can be chosen and $n!$ ways in which arrows can be drawn from P to Q . Each combination produces a distinct nice partition, so the number of nice partitions is $2n \times n!$. \square

2.7

A 2002×2002 square is divided into unit squares. A straight line may be drawn, and every unit square whose interior it intersects is colored. Prove that at least 1173 lines are required to color the whole square.

Solution: Let $n = 2002$. We index the unit squares with double coordinates, with $(1, 1)$ being in the bottom-left corner. We shall call “left chain” the sequence of squares

$$(i_1, j_1), (i_2, j_2), \dots, (i_{2n-1}, j_{2n-1}),$$

where $(i_1, j_1) = (1, 1)$ and either $(i_{k+1}, j_{k+1}) \equiv (i_k + 1, j_k)$ or $(i_{k+1}, j_{k+1}) \equiv (i_k, j_k + 1)$ for $k = 1, 2, \dots, 2n - 2$. Similarly, a “right chain” is defined as a sequence that starts at $(i_1, j_1) = (1, n)$ and either $(i_{k+1}, j_{k+1}) \equiv (i_k + 1, j_k)$ or $(i_{k+1}, j_{k+1}) \equiv (i_k, j_k - 1)$ for $k = 1, 2, \dots, 2n - 2$.

Observe that the squares intersected by any line form a subset of either a left chain or a right chain. Indeed, consider the segment in which the line intersects the big square. Traverse this segment from its lower endpoint to its upper endpoint. As we move along the segment, the y -coordinate never decreases. If the slope of the line is nonnegative, then the x -coordinate also never decreases. Consequently, each new square intersected by the segment lies either to the right of the previous one, above it, or diagonally above-right (when the segment passes through a grid vertex). Thus the colored squares appear in a monotone order and can be embedded in a left chain, although some squares of the chain may be skipped if the segment passes through a vertex. If the slope of the line is negative, then x never increases while y never decreases, and the same reasoning shows that the colored squares form a subset of a right chain.

We shall prove that at least $(2 - \sqrt{2})n$ chains are needed to cover the big square.

Let us assume the contrary, and suppose that there exist s chains, $s < (2 - \sqrt{2})n$ that cover the big square. We order the chains arbitrarily and take the first one. In each of its squares, we write its ordinal number in the chain, i.e. in the square (i_m, j_m) we write m . Then we take the second chain and, again, write in each of its squares its ordinal number in the chain, unless it already contains a number written by a previous chain. We proceed the same way for all chains, so after s steps each square will have a number.

Let $c(m)$ denote the number of times that the number m is written in the entire table. We note that each square (i, j) contains either the number $i + j - 1$, if it was written by a left chain, or the number $i + n - j$, if it was written by a right chain, so for m to be written in a square, the pair (i, j) must satisfy one of the equations $i + j - 1 = m$ or $i + n - j = m$.

For $m \leq n$, there are $2m$ possible solutions to the above equations, and they are given by the diagonals $(1, m), (2, m - 1), \dots, (m, 1)$, and $(m, n), (m - 1, n - 1), \dots, (1, n - m + 1)$, so $c(m) \leq 2m \leq 2(2n - m)$.

Similarly, for $m > n$, there are $2(2n - m)$ possible solutions given by the diagonals $(m - n, n), (m - n + 1, n - 1), \dots, (n, m - n)$, and $(m - n + 1, 1), (m - n + 2, 2), \dots, (n, 2n - m)$, so $c(m) \leq 2(2n - m) \leq 2m$.

Therefore, in all cases we have

$$c(m) \leq 2m \text{ and } c(m) \leq 2(2n - m).$$

On the other hand, $c(m) \leq s$, since the number m can occur in a given chain at most once. Combining with the earlier upper bound on $c(m)$, we get

$$c(m) \leq \min(2m, 2(2n - m), s).$$

Since the maximum possible count of occurrences depends on whether s is even or odd, consider the two cases separately.

Case 1, $s = 2p$. Then the numbers

$$1, 2, \dots, p, p+1, \dots, n, n+1, \dots, 2n-p, 2n-p+1, \dots, 2n-1$$

occur respectively at most

$$2, 4, \dots, 2p, 2p, \dots, 2p, 2p, \dots, 2p, 2p-2, \dots, 2$$

times. We have at most

$$2(1+2+\dots+p) + (2n-1-2p)2p + 2(1+2+\dots+p) = 2p(p+1) + (2n-1)2p - p^2 = 4np - 2p^2$$

numbered squares, hence $n^2 \leq 4np - 2p^2$. That gives us

$$p \geq \left(1 - \frac{\sqrt{2}}{2}\right)n,$$

whence $s = 2p \geq (2 - \sqrt{2})n > 1172$.

Case 2, $s = 2p - 1$. Then the numbers

$$1, 2, \dots, p-1, p, \dots, n, n+1, \dots, 2n-p, \dots, 2n-1$$

occur respectively at most

$$2, 4, \dots, 2p-2, s, \dots, s, 2p-2, \dots, 2$$

times. We have at most

$$2(1+2+\dots+p-1) + (2n-1-2(p-1))s + 2(1+2+\dots+p-1) = 2p(p-1) + (2n-s)s$$

numbered squares, hence

$$n^2 \leq (s-1)(s+1)/2 + 2ns - s^2,$$

which gives us $s \geq 2n - \sqrt{2n^2 - 1} > 1172$. In either case, we have $s \geq 1173$. \square

2.8

There are $2n$ cards arranged in a row. The numbers $1, 2, \dots, 2n$ are written on the two sides of the cards, each number appearing exactly twice in total. Initially only one side of each card is visible.

A move consists of flipping one card. The goal is to reach a configuration in which all visible numbers are distinct.

- (i) Prove that $3n - 2$ moves always suffice.
- (ii) Prove that $3n - 3$ moves do not always suffice.

Solution: Without loss of generality, we can treat the numbers as simply any $2n$ distinct numbers.

(i) We shall prove the statement by induction. For $n = 1$, either the two numbers shown are equal, so we turn either card, or they are different and there is nothing to do. So $3n - 2 = 1$ moves suffice.

Now, assume that the statement holds for $1, 2, \dots, n - 1$. If all shown numbers are different, we are done. If they are not, assume without loss of generality that the number appearing twice is 1. Let those two cards be called “starting” and the remaining $2n - 2$ cards be called “remaining”. We flip both starting cards and let the uncovered numbers be a and b .

Define $N = \{1, 2, \dots, 2n\}$.

- If $a = b$, then $a, b \neq 1$, so the numbers 1 and a are used twice across the starting cards and the other $2n - 2$ numbers $N \setminus \{1, a\}$ are used twice across the remaining $2n - 2$ cards. That means the induction hypothesis applies to the remaining cards and we need at most $3(n - 1) - 2$ moves to display all numbers in $N \setminus \{1, a\}$ in the remaining cards. In the end, we flip back one of the starting cards to uncover the 1 on its back, and we are done with a total of $2 + 3(n - 1) - 2 + 1 = 3n - 2$ moves.
- If $a \neq b$, then each of the numbers a and b is used once more in the remaining cards. We temporarily relabel those two occurrences of a and b in the remaining cards with some arbitrary number $x \notin N$. The $2n - 2$ remaining cards now contain $2n - 2$ distinct numbers $N \setminus \{1, x\}$, each used twice, so the induction hypothesis applies and we need at most $3(n - 1) - 2$ moves to display all those numbers exactly once. The x that now appears was originally either a or b , so depending on which one it was, we flip back one of the starting cards to uncover the 1 on its back. Again, we are done with a total of $2 + 3(n - 1) - 2 + 1 = 3n - 2$ moves.

(ii) Consider a starting configuration where the numbers $1, 2, \dots, n$ appear twice on the front and the numbers $n + 1, n + 2, \dots, 2n$ are written twice on the back of the cards. No matter what strategy we choose, sooner or later there must be a move that uncovers the number $2n$. At any move, we shall call a card “primary” if it has not been flipped yet and “secondary” if it has been flipped already.

Notice that (a) swapping two moves does not change the resulting configuration; and (b) flipping a secondary card carries no new information, since we know what is on its

back. Combining (a) and (b), we can assume that any move flipping a secondary card is postponed until after there are no more primary cards.

We know that the number $2n$ is written twice on the back of two cards, but we do not know which ones those are. In the worst case, the two cards containing $2n$ may be the last two primary cards flipped. Thus it may take $2n - 1$ moves before the first $2n$ is revealed. Let i , $1 \leq i \leq n$ be the number on the front of the last remaining primary card..

Each of the $2n - 2$ cards originally displaying

$$1, 1, 2, 2, \dots, i - 1, i - 1, i + 1, i + 1, \dots, n, n$$

is now flipped, so from each pair of cards with j written on their back, $1 \leq j \leq n, j \neq i$, at least one must be flipped again. That means we need at least a total of $(2n - 1) + (n - 1) = 3n - 2$ moves, which completes the proof of (ii). \square

3 Number Theory

3.1

Let p be prime. There are p numbers written in a circle. At each move, simultaneously each number is replaced by the number plus its right neighbor minus twice its left neighbor. Prove that after $p - 1$ moves all numbers will give equal remainders upon division by p .

Solution: We shall consider the numbers modulo p . Define two operators $d(x)$ and $f(x)$ on any vector $x = (x_1, x_2, \dots, x_p)$ as follows:

$$d(x) = (x_1 - x_2, x_2 - x_3, \dots, x_p - x_1),$$

$$f(x) = (x_1 + x_2 - 2x_p, x_2 + x_3 - 2x_1, \dots, x_p + x_1 - 2x_{p-1}).$$

The operator f corresponds exactly to the move applied to the vector of the p numbers.

Define $\vec{0} = (0, 0, \dots, 0)$ and call a vector *uniform* if all its entries are the same. We shall list the following properties of d and f :

- For any uniform vector x , $f(x) = \vec{0}$ and $d(x) = \vec{0}$. (*)
- If $d(x) = \vec{0}$, then $x_1 - x_2 = x_2 - x_3 = \dots = 0$, so all x_i are the same and hence x is uniform. (**)
- Direct computation shows that

$$d(f(x)) = (3x_1 - 2x_p - x_3, 3x_2 - 2x_1 - x_4, \dots, 3x_p - 2x_{p-1} - x_2),$$

which coincides with $f(d(x))$. Hence d and f commute. (***)

Next, we show by induction that

$$d^k(x) = \left(\sum_{i=0}^k \binom{k}{i} (-1)^i x_{1+i}, \sum_{i=0}^k \binom{k}{i} (-1)^i x_{2+i}, \dots, \sum_{i=0}^k \binom{k}{i} (-1)^i x_{p+i} \right),$$

where the indices are taken cyclically modulo p (i.e. $x_{p+1} = x_1$, and so on). The statement is obvious for $k = 1$. If we assume that it holds for k , then

$$d^{k+1}(x) = d \left(\sum_{i=0}^k \binom{k}{i} (-1)^i x_{1+i}, \sum_{i=0}^k \binom{k}{i} (-1)^i x_{2+i}, \dots, \sum_{i=0}^k \binom{k}{i} (-1)^i x_{p+i} \right).$$

The r -th term of $d^{k+1}(x)$ is then

$$\begin{aligned} & \sum_{i=0}^k \binom{k}{i} (-1)^i x_{r+i} - \sum_{i=0}^k \binom{k}{i} (-1)^i x_{r+i+1} \\ &= \binom{k}{0} (-1)^0 x_{r+0} + \sum_{i=1}^k \left(\binom{k}{i} (-1)^i - \binom{k}{i-1} (-1)^{i-1} \right) x_{r+i} - \binom{k}{k} (-1)^k x_{r+k+1} \end{aligned}$$

$$\begin{aligned}
&= \binom{k}{0} x_r + \sum_{i=1}^k \left(\binom{k}{i} + \binom{k}{i-1} \right) (-1)^i x_{r+i} + \binom{k+1}{k+1} (-1)^{k+1} x_{r+k+1} \\
&= \sum_{i=0}^{k+1} \binom{k+1}{i} (-1)^i x_{r+i},
\end{aligned}$$

and the induction step is complete. Applying the formula with $k = p$ and working modulo p , since $p \mid \binom{p}{i}$ for $0 < i < p$, we have

$$d^p(x) = (x_1 - x_1, x_2 - x_2, \dots, x_p - x_p) = \vec{0}.$$

From (**) we have that $d^{p-1}(x)$ is uniform, so using (*) we have $f(d^{p-1}(x)) = \vec{0}$. Using (***) , that gives us $d^{p-1}(f(x)) = \vec{0}$.

Again, from (**) we have that $d^{p-2}(f(x))$ is uniform, so using (*) we have $f(d^{p-2}(f(x))) = \vec{0}$. Using (***) , that gives us $d^{p-2}(f^2(x)) = \vec{0}$.

Continuing this argument inductively, we obtain that $d^{p-1-i}(f^i(x))$ is uniform for $i = 0, 1, \dots, p-1$. Using $i = p-1$ and setting x to the vector of the numbers on the circle, we get that $f^{p-1}(x)$ is uniform, which completes the proof. \square

3.2

Let d be the smallest positive integer for which $n \mid 2^d + 1$. If $n > 6d$, prove that there are more than $6d$ positive integers m , $m \leq n$, such that for some $x, y, z \in \mathbb{N}$,

$$n \mid (2^x + 2^y + 2^z + m).$$

Solution: From $n \mid 2^d + 1$ it follows that n is odd and greater than 6. We shall consider the numbers as residues modulo n .

Let X, Y, Z be the sets of all *nonzero* residues modulo n representable as 2^x , $2^x + 2^y$, and $2^x + 2^y + 2^z$ respectively. Notice that $X \subseteq Y$, since $2^x \equiv 2^{x+2d-1} + 2^{x+2d-1} \in Y$ and that $Y \subseteq Z$, since $2^x + 2^y \equiv 2^{x+2d-1} + 2^{x+2d-1} + 2^y \in Z$. In view of the fact that $2^d + 2^d + 2^1 + n \equiv 0$, we see that n is an admissible m , and since $n \equiv 0 \notin Z$, the remaining admissible values of m correspond to the nonzero residues in Z . Thus it suffices to prove that $|Z| \geq 6d$.

Since $2^{2d} \equiv (-1)^2 = 1$, there exists a *minimal* j such that $2^j \equiv 1$ and $j \mid 2d$. Notice that $j \neq d$, otherwise $1 \equiv -1$, which is impossible for $n > 6$. At the same time, it cannot be that $j < d$, otherwise $2^{d-j} \equiv -1$ would contradict the minimality of d . Therefore $j > d$.

Since $j > d$ and $j \mid 2d$, we must have $j = 2d$. Thus $X = \{2^i\}, i = 1, 2, \dots, 2d$, so $|X| = 2d$.

Consider first the case where $X = Y$. In that case, for every $a \in X$, from $2^{2d} \equiv 1$ we have $a + 1 \equiv a + 2^{2d} \in Y = X$, hence recursively we get $a, a + 1, a + 2, \dots, n - 1 \in X$. From $n - 1 + 2^1 \in Y = X$ we get that $n + 1 \equiv 1 \in X$, and again recursively we get that $2, 3, \dots, a - 1 \in X$. So X contains all $n - 1$ nonzero residues. Therefore $|X| = n - 1$, and so $|Z| \geq |X| = 6d$ and the condition is proved.

So from now on, we assume that $X \subset Y$. We choose an arbitrary $a \in Y$. Since n is odd and $a \not\equiv 0$, we have $2a \not\equiv 0$ and so $2a \in Y$. Consider a bijection $f : Y \rightarrow Y$ defined by $f(a) = 2a$. It is an injection, since $2a \equiv 2b \Rightarrow a = b$, and hence it is a bijection, since it is an injection of one set into itself. For any $a \in Y$, consider its orbit in f of length k , so k is the minimal number such that $f^k(a) = a$. We have $a2^k \equiv a \Rightarrow 2^k \equiv 1$, since $a \not\equiv 0$. By the minimality of $2d$ we get $2d \mid k$, and from $k \leq 2d$ we get $k = 2d$. That means all orbits are of length $2d$, hence $2d \mid |Y|$.

Similarly to the earlier case $X = Y$, if $Y = Z$ we end up with $|Y| = n - 1$ and $|Z| \geq 6d$, so we can consider $Y \subset Z$. Like before, we conclude $2d \mid |Z|$.

Therefore we have that

$$\frac{|X|}{2d} < \frac{|Y|}{2d} < \frac{|Z|}{2d},$$

are three distinct integers. So $\frac{|Z|}{2d} \geq 3$ and $|Z| \geq 6d$, which completes the proof. \square

3.3

Let p be prime and $a \neq b$ be positive integers less than p . Prove that the equation

$$am + bn = p$$

has a solution in positive integers m, n if and only if the sets

$$A = \left\{ \left\lfloor \frac{ip}{a} \right\rfloor : i = 1, 2, \dots, a-1 \right\}, \quad B = \left\{ \left\lfloor \frac{jp}{b} \right\rfloor : j = 1, 2, \dots, b-1 \right\}$$

have no common element.

Solution: We shall prove each direction in turn.

Forward direction. We have $am + bn = p$ and want to show that $A \cap B = \emptyset$. Suppose the contrary, and let k be an element of $A \cap B$, so

$$k = \left\lfloor \frac{ip}{a} \right\rfloor = \left\lfloor \frac{jp}{b} \right\rfloor \quad (*)$$

for some $1 \leq i \leq a-1$ and $1 \leq j \leq b-1$. That gives us

$$\frac{ip}{a} > k > \frac{ip}{a} - 1 \quad \text{and} \quad \frac{jp}{b} > k > \frac{jp}{b} - 1,$$

where the strictness of the inequalities comes from the fact that the elements inside the floor brackets in (*) are non-integer. Multiply the first inequality by am and the second by bn , then add them to obtain

$$ipm + jpn > kam + kbn > ipm - am + jpn - bn.$$

Using $am + bn = p$ and dividing by p , we get

$$im + jn > k > im + jn - 1,$$

which is impossible. So $k \in A \cap B$ cannot exist and we are done.

Backward direction. We have $A \cap B = \emptyset$ and want to prove that $am + bn = p$ for some positive integers m, n . If $d = \gcd(a, b) > 1$, we have $a = a_1d$ and $b = b_1d$ for some $a_1 < a$ and $b_1 < b$, hence

$$\left\lfloor \frac{a_1p}{a} \right\rfloor = \left\lfloor \frac{p}{d} \right\rfloor = \left\lfloor \frac{b_1p}{b} \right\rfloor,$$

and so $\left\lfloor \frac{p}{d} \right\rfloor \in A \cap B$, contradicting $A \cap B = \emptyset$. Therefore, $d = 1$, so there exists $m' \in \{1, 2, \dots, b-1\}$ such that $am' \equiv 1 \pmod{b}$. Writing $am' - 1 = n'b$ yields

$$m'a - n'b = 1$$

with $n' \in \{1, 2, \dots, a-1\}$. Next,

$$\frac{m'p}{b} - \frac{n'p}{a} = \frac{p}{ab} > 0 \Rightarrow \left\lfloor \frac{m'p}{b} \right\rfloor \geq \left\lfloor \frac{n'p}{a} \right\rfloor,$$

where equality is not possible, otherwise A and B would have a common element, contradicting $A \cap B \neq \emptyset$. Thus

$$\left\lfloor \frac{m'p}{b} \right\rfloor = \left\lfloor \frac{n'p}{a} \right\rfloor + k$$

for some integer $k \geq 1$. Therefore,

$$\frac{p}{ab} = \frac{m'p}{b} - \frac{n'p}{a} = \left\lfloor \frac{m'p}{b} \right\rfloor - \left\lfloor \frac{n'p}{a} \right\rfloor + \left\{ \frac{m'p}{b} \right\} - \left\{ \frac{n'p}{a} \right\} = k + \left\{ \frac{m'p}{b} \right\} - \left\{ \frac{n'p}{a} \right\}.$$

Multiplying both sides by ab and rearranging, we get

$$p = \left\{ \frac{m'p}{b} \right\} ab + \left(k - \left\{ \frac{n'p}{a} \right\} \right) ab = am + bn,$$

where

$$m = b \left\{ \frac{m'p}{b} \right\} = m'p - b \left\lfloor \frac{m'p}{b} \right\rfloor$$

and

$$n = a \left(k - \left\{ \frac{n'p}{a} \right\} \right) = ak - n'p + a \left\lfloor \frac{n'p}{a} \right\rfloor$$

are positive integers, so the statement is proved. □

3.4

For which positive integers n can the divisors of 2001^n be partitioned into triples such that the product of the numbers in each triple is the same?

Solution: Observe that $2001 = pqr$ for the primes $p = 3, q = 23, r = 29$. Since the total number of divisors of $(pqr)^n$ is $(n + 1)^3$, the number of triples is $(n + 1)^3/3$ and so $3 \mid (n + 1)$. Let the product of the numbers in each triple be a . If we multiply all these triples, we obtain

$$a^{(n+1)^3/3} = (pqr)^{(n+1)^2(0+1+\dots+n)} = (pqr)^{(n+1)^3 n/2},$$

hence $a = 2001^{3n/2}$, so n is even. Combining with $3 \mid n + 1$, a necessary condition for n becomes $n \equiv 2 \pmod{6}$. We shall prove that this condition is also sufficient.

Let $n = 6m + 2$ for some integer $m \geq 0$, and let A denote the set of $2m + 1$ vectors α_k defined as

$$\alpha_k = (k, m + k, 2m - 2k), \quad k = 0, 1, \dots, m$$

and

$$\alpha_k = (k, k - m - 1, 4m + 1 - 2k), \quad k = m + 1, \dots, 2m.$$

For each α , observe that the sum of its coordinates is $3m$. Furthermore, observe that each coordinate assumes every value $0, 1, \dots, 2m$ exactly once as k iterates from 0 to $2m$. Indeed, the first coordinate takes the values $0, 1, \dots, m, m+1, \dots, 2m$; the second coordinate takes the values $m, m+1, \dots, 2m, 0, 1, \dots, m-1$; and the third coordinate takes the values $2m, 2m-2, \dots, 0, 2m-1, 2m-3, \dots, 1$.

Define the transformation f by

$$f(x, y, z) = (3x, 3y + 1, 3z + 2)$$

and the set F as:

$$F = \bigcup_{\alpha \in A} f(\alpha).$$

Notice that F has the following properties:

- (i) all coordinates of its vectors are in the interval $[0, n]$, since $x, y, z \in [0, 2m]$;
- (ii) every integer in $\{0, 1, \dots, n\}$ appears as one of the coordinates in exactly one vector from F , since $3x$ iterates over all numbers congruent to $0 \pmod{3}$, $3y + 1$ over all numbers congruent to $1 \pmod{3}$, and $3z + 2$ over all numbers congruent to $2 \pmod{3}$;
- (iii) the sum of coordinates of each vector in F is exactly $3(x + y + z) + 3 = 9m + 3$.

Similarly, define

$$f'(x, y, z) = (3x + 2, 3y, 3z + 1), \quad f''(x, y, z) = (3x + 1, 3y + 2, 3z)$$

and

$$F' = \bigcup_{\alpha \in A} f'(\alpha), \quad F'' = \bigcup_{\alpha \in A} f''(\alpha).$$

Analogously, properties (i), (ii), and (iii) apply to F' and F'' . Furthermore, the sets F, F', F'' are disjoint, since the first coordinates have different residues modulo 3. Let $U = F \cup F' \cup F''$. Note that by property (ii), every integer in $\{0, 1, \dots, n\}$ appears exactly once as a coordinate of a vector in F , and exactly three times among the vectors in U , once in each coordinate position.

We construct the partitioning of the triples as follows: for every element

$$(x_1, y_1, z_1) \in F, \quad (x_2, y_2, z_2), (x_3, y_3, z_3) \in U,$$

define the triple

$$(p^{x_1} q^{x_2} r^{x_3}, p^{y_1} q^{y_2} r^{y_3}, p^{z_1} q^{z_2} r^{z_3}).$$

For every given triple, from (i), each number is a divisor of $p^n q^n r^n = 2001^n$, and from (iii), the product of these numbers is the constant

$$p^{x_1+y_1+z_1} q^{x_2+y_2+z_2} r^{x_3+y_3+z_3} = (pqr)^{9m+3}.$$

Furthermore, the numbers in each triple are different, since x_1, y_1, z_1 are congruent to 0, 1, 2 (mod 3), the exponent vectors differ in at least one coordinate.

What remains is to show that this set of triples constitutes an actual *partition* of the divisors of 2001^n . Consider any one of these divisors $p^{u_1} q^{u_2} r^{u_3}$. We need to show that it appears in exactly one of the triples.

From (ii), we know that u_1 appears exactly once as a coordinate of exactly one vector of F . Let that vector be $\mu_1 = (x_1, y_1, z_1)$, and let $l \in \{1, 2, 3\}$ denote the coordinate position of u_1 in it.

Again from (ii), among the vectors in U , there are exactly 3 vectors that contain u_2 as a coordinate, each in a different position. Let the one that has u_2 in position l be $\mu_2 = (x_2, y_2, z_2)$. Similarly, $\mu_3 = (x_3, y_3, z_3)$ is determined. Notice that the triple corresponding to $\mu_1 \in F, \mu_2, \mu_3 \in U$ contains $p^{u_1} q^{u_2} r^{u_3}$ in position l . That completes the proof.

Finally, all desired n are given by $n = 2 + 6m, \quad m = 0, 1, \dots$ □

3.5

Let $p > 2$ be prime, and let \underline{x} denote the least non-negative residue of x modulo p . For $0 \leq a < p$ define

$$f_a(m) = m + \underline{ma}.$$

Determine the number of integers a with $0 \leq a < p$ such that

$$f_a(m) > a \quad \text{for all } m = 1, 2, \dots, p-1.$$

Solution: We divide the proof into three parts. The first proposes a set of numbers that work; the second shows that those are the only numbers that work; and the third part counts them. Define $a_k = \left\lfloor \frac{p}{k} \right\rfloor$ for all $k \geq 1$ and let

$$A = \{a_k \mid k = 2, 3, \dots, p\}.$$

We call a number $a \in [0, p-1]$ *good* if $f_a(m) > a$ for all $m \in [1, p-1]$. Our goal is to count the good numbers.

Lemma 1. We shall prove that every $a \in A \cup \{0, p-1\}$ is a good number.

Proof: If $a = 0$, then $f_a(m) = m > 0 = a$; and if $a = p-1$, then $f_a(m) = m + \underline{ma} = m + \underline{-m} = m + p - m = p > p-1 = a$. So 0 and $p-1$ are good numbers. Next, let $a \in A$. We shall prove by contradiction that there cannot exist $m_0 \in [1, p-1]$ such that $a \geq f_a(m_0)$. Assume the contrary, and let m_0 be the number minimizing $f_a(m)$, i.e.

$$a \geq f_a(m_0) = \min_{1 \leq m \leq p-1} f_a(m).$$

From $f_a(1) = 1 + a > a$, we see that $m_0 \neq 1$, so $a \geq f_a(m_0) \geq m_0 > 1$, thus $a > 1$. Let $v = \underline{m_0 a} > 0$ and $m_0 a = lp + v$ for some $l \geq 0$. From $m_0 > 1$ and the minimality of m_0 , we have

$$m_0 + v = f_a(m_0) \leq f_a(m_0 - 1) = (m_0 - 1) + \underline{(m_0 - 1)a},$$

so $v < \underline{(m_0 - 1)a}$. It must be that $v < a$, otherwise $\underline{(m_0 - 1)a} = \underline{v - a} = v - a \leq 0$, which would contradict $v < \underline{(m_0 - 1)a}$. Hence $0 \leq v < a$, and thus

$$\frac{lp}{a} < m_0 < \frac{lp}{a} + 1 \Rightarrow m_0 = \left\lceil \frac{lp}{a} \right\rceil.$$

Since $a \in A$, there exist integers $k, k' : 0 < k' < k$ for which $a = a_k$ and $p = ak + k'$, so

$$m_0 = \left\lceil \frac{lp}{a} \right\rceil = \left\lceil \frac{l(ak + k')}{a} \right\rceil = lk + \left\lceil \frac{lk'}{a} \right\rceil.$$

From $a \geq f_a(m_0) > m_0 \geq lk > lk'$ we get

$$\left\lceil \frac{lk'}{a} \right\rceil = 1 \Rightarrow m_0 = lk + 1.$$

Using again $a > lk'$ and $ak = p - k'$, we have

$$f_a(m_0) = f_a(lk + 1) = lk + 1 + \underline{lka + a} = lk + 1 + \underline{a - lk'}.$$

But as $0 \leq a - lk' < a < p$, we have $\underline{a - lk'} = a - lk'$, so

$$f_a(m_0) = lk + 1 + (a - lk') = (a + 1) + l(k - k') \geq a + 1 > a,$$

which contradicts the choice of m_0 , and Lemma 1 is proved.

Lemma 2. We shall prove that if $a \notin A$ and $0 < a < p - 1$, then $f_a(m) \leq a$ for some m .

Proof: Since $a \notin A$ and $a_p = 1 < a < p = a_1$, there must exist $k \in [1, p - 1]$ such that

$$a_{k+1} < a < a_k.$$

For $p > 2$ we have $a_{p-1} = a_p = 1$, so $k \neq p - 1$ and hence $k \in [1, p - 2]$. From the upper bound, we get

$$a + 1 \leq a_k \leq \frac{p}{k},$$

where at least one of the two inequalities is strict, except when $k = 1$ and $a = p - 1$, which contradicts $a < p - 1$. From the lower bound, we have

$$a \geq a_{k+1} > \frac{p}{k+1}.$$

From $p < ak + a < p + a < 2p$ we have $\underline{ak + a} = (ak + a) - p$, so

$$ak + k < p < ak + a \Rightarrow f_a(k + 1) = k + 1 + \underline{ak + a} = k + 1 + ak + a - p < a + 1,$$

thus $f_a(k + 1) \leq a$ and Lemma 2 is proved.

Lemmas 1 and 2 tell us that all the good numbers are $0, p - 1$ and the elements of A . All it remains then is to count $|A|$ and add 2. Intuitively, for small k , the numbers a_k are widely spaced, so they are all different; and for large k , a_k are thinly spaced, so they assume every integer value from some point onward.

Let k_0 be the largest $k \in [2, p - 1]$ for which $\frac{p}{k-1} - \frac{p}{k} > 1$. Such a number necessarily exists, since $k = 2$ satisfies the inequality for $p > 2$. If the gap between two rational numbers is > 1 , their floors differ by at least 1. Applying this observation to our case yields $a_k < a_{k-1}$ for $2 \leq k \leq k_0$, hence a_2, a_3, \dots, a_{k_0} are $k_0 - 1$ distinct elements of A . On the other hand, when the gap between two rational numbers is ≤ 1 , their floors differ by at most 1. Applying this to our case, for $k > k_0$ we have $a_{k-1} - a_k \leq 1$, so all positive numbers less than a_{k_0} also belong to A . Therefore

$$|A| = k_0 - 1 + a_{k_0} - 1 = k_0 + \left\lfloor \frac{p}{k_0} \right\rfloor - 2.$$

We just need to find the k_0 and compute the above expression. So k_0 must satisfy the inequalities:

$$\frac{p}{k-1} - \frac{p}{k} > 1 \geq \frac{p}{k} - \frac{p}{k+1}$$

$$\Rightarrow k(k+1) > p > k(k-1).$$

Notice that $s = \sqrt{p}$ is irrational. Consider two cases:

Case 1. $p > k^2$, so $k(k+1) > p > k^2$. We have

$$(2k+1)^2 > 4(k+1)k > 4p > 4k^2 \Rightarrow s \in (k, k+1/2),$$

so $k = \lfloor s \rfloor$. From $k+1 > p/k > k$ we obtain $a_k = k$, and hence $|A| = a_k + k - 2 = 2k - 2 = 2\lfloor s \rfloor - 2 = \lfloor 2s \rfloor - 2$.

Case 2. $p < k^2$, so $k^2 > p \geq k(k-1) + 1$. We have

$$4k^2 > 4p \geq 4k(k-1) + 4 > (2k-1)^2 \Rightarrow s \in (k-1/2, k),$$

so $k = \lfloor s \rfloor + 1$. From $k > p/k > k-1$ we obtain $a_k = k-1$, and so $|A| = a_k + k - 2 = 2k - 3 = 2\lfloor s \rfloor - 2 = \lfloor 2s \rfloor - 2$.

The case $p = k^2$ need not be examined, since s is irrational. Therefore $|A| = \lfloor 2\sqrt{p} \rfloor - 2$, and adding back the 2 good numbers 0 and $p-1$, the final count becomes $\lfloor 2\sqrt{p} \rfloor$. \square

3.6

Let p be a prime and consider a regular $2p$ -gon. Let A and B be opposite vertices. Among the remaining vertices, $k < p$ are marked. Draw all segments joining pairs of marked vertices that intersect AB , and assume that at least one such segment exists. Prove that at least $k - 1$ of these segments can be chosen so that no two are parallel or perpendicular.

Solution: Number the p vertices from A to B along one side with $0, 1, \dots, p - 1$ and call this set L . Likewise, number the p vertices from B to A on the other side and call this set R . Every segment crosses AB , so it must have one endpoint numbered $a \in L$ and the other $b \in R$ due to the convexity of the polygon. Since vertex k corresponds to angle $k\pi/p$, the direction of chord ab depends only on the midpoint angle $(a + b)\pi/(2p)$. Hence two chords are parallel iff these midpoint angles are equal, and perpendicular iff they differ by $\pi/2$. Therefore,

$$\begin{aligned} a_1b_1 \parallel a_2b_2 &\Leftrightarrow a_1 + b_1 = a_2 + b_2, \\ a_1b_1 \perp a_2b_2 &\Leftrightarrow a_1 + b_1 = a_2 + b_2 \pm p. \end{aligned}$$

In either case, the statement that a_1b_1 is either parallel or perpendicular to a_2b_2 is equivalent to $a_1 + b_1 \equiv a_2 + b_2 \pmod{p}$. So all we need to do is prove the following Lemma:

Lemma. Let A, B be sets of residues modulo p and $|A| + |B| < p$. Define their “sum” by

$$A + B = \{a + b \mid a \in A, b \in B\}.$$

Then

$$|A + B| \geq |A| + |B| - 1.$$

Proof: Let $|A| = m$, $|B| = n$, and assume without loss of generality that $m \leq n$. Define “translation” t on the set C as follows:

$$t + C = \{t + c \mid c \in C\}.$$

Observe that $(t + A) + B = t + (A + B)$, hence $|(t + A) + B| = |A + B|$. If $m = 1$, then $|A + B| = |B| = |A| + |B| - 1$ and we are done, so hereafter assume $m > 1$ and let a_1, a_2 be two elements of A . Furthermore, $m \leq n < p$, hence $\exists c : c \notin B$. Now, as the sequence

$$c + t(a_2 - a_1), \quad t = 1, 2, \dots, p - 1$$

runs through all residues modulo p except c , there exists a minimal t such that for $b = c + t(a_2 - a_1) \in B$. Then

$$A' = (b - a_2) + A$$

contains the element $b + (a_1 - a_2) \in B$ and $b + (a_2 - a_2) = b \notin B$. Since A' is just a translation of A , we only need to prove the lemma for A' and B . Let

$$F = A' \cap B, \quad G = A' \cup B.$$

We have

$$\emptyset \neq F \subset A', \quad B \subset G,$$

hence

$$0 < |F| < m, n < |G|.$$

Observe that $|F| + |G| = |A| + |B|$, since every element $f \in F$ is counted exactly twice on each side, and every element $x \in (A' \cup B) \setminus F$ is counted exactly once on each side. We shall show that

$$F + G \subseteq A + B.$$

Indeed, if $f \in F, g \in G$, then depending on whether $g \in A'$ or $g \in B$ we have $(f, g) \in A' \times B$ or $(f, g) \in B \times A'$. In either case, $f + g \in A' + B$. Thus

$$|A' + B| \geq |F + G|,$$

so it suffices to prove the lemma for F and G instead of A' and B . Thus we replace the pair A, B (with $|A| \leq |B|$) in the original statement by the pair F, G , for which $|F| < |A|$ and $|G| > |B|$. Repeating this reduction decreases the size of the smaller set each time, so eventually that set has size 1, which was already handled. Thus the lemma is proved. Applying the lemma to the original problem, each segment corresponds to the residue $a + b \pmod{p}$. Since parallel or perpendicular segments correspond to the same residue, selecting segments with distinct residues yields a family of at least $k - 1$ segments no two of which are parallel or perpendicular. \square

4 Geometry

4.1

Let $ABCD$ be an isosceles trapezoid with $AB \parallel CD$. A point T is taken on AB such that $DP \cdot DT = CQ \cdot CT$, where $P = AC \cap TD$ and $Q = BD \cap TC$. Prove that the circumcenter of $\triangle TPQ$ is equidistant from A and B .

Solution: Let $\angle DCA = \alpha$, $\angle DCT = y$, $\angle CDT = x$. Then we have:


$$\begin{aligned} \frac{CD \sin \alpha}{\sin(x + \alpha)} \cdot \frac{CD \sin y}{\sin(x + y)} &= DP \cdot DT \\ &= CQ \cdot CT = \frac{CD \sin \alpha}{\sin(y + \alpha)} \cdot \frac{CD \sin x}{\sin(x + y)} \\ \Rightarrow \frac{\sin y}{\sin(x + \alpha)} &= \frac{\sin x}{\sin(y + \alpha)} \end{aligned}$$

Therefore

$$\begin{aligned} \sin y(\sin y \cos \alpha + \sin \alpha \cos y) &= \sin x(\sin x \cos \alpha + \sin \alpha \cos x) \\ \Rightarrow \cos \alpha(\sin^2 y - \sin^2 x) &= \sin \alpha(\sin x \cos x - \sin y \cos y) \\ \Rightarrow \cos \alpha \sin(x + y) \sin(y - x) &= \sin \alpha \cos(x + y) \sin(y - x) \\ \Rightarrow \sin(x + y + \alpha) \sin(x - y) &= 0 \\ \Rightarrow x = y \quad \text{or} \quad x + y + \alpha &= 180^\circ \end{aligned}$$

In the first case, $\triangle DTC$ is isosceles and symmetric with respect to the perpendicular bisector of AB and of DC (hereafter called just *symmetric* for simplicity). So T is the midpoint of AB , and P and Q are symmetric, thus the circumcenter of $\triangle TPQ$ is on the line of symmetry, and we are done.

In the second case, $\angle DTC = 180^\circ - x - y = \alpha$. Let T' be the point symmetric to T , $P' = AC \cap T'D$, and $Q' = BD \cap T'C$. Notice that P' and Q' are symmetric to Q and P respectively, so $PQ' \parallel DC$ and therefore $\angle PQ'Q = \alpha = \angle PTQ$. That means PQ is seen at the same angle from Q' and T and so $TPQQ'$ are concyclic. Hence point T (and therefore also T') lies on the circumcircle about the isosceles trapezoid $PQQ'P'$. Again due to symmetry, the circumcenter of $\triangle TPQ$ is on the line of symmetry and so equidistant from A and B . \square



4.2

From the center of the circle k a perpendicular is dropped to the line l , intersecting k at the point D . Let P, Q be arbitrary points of l , and

$$P' = (PD) \cap k, \quad Q' = (QD) \cap k, \quad P, Q \neq D.$$

Prove that the center of the circumcircle of $\triangle P'Q'P$ lies on k exactly when

$$PQ = P'Q'.$$

Solution: Obviously l is external to k . An inversion with center D which maps k onto l (and vice versa) sends P into P' and Q into Q' , hence P, P', Q, Q' lie on a circle.

Let the positive direction of rotation about D be counterclockwise. If

$$H = (OD) \cap l,$$

we define

$$OH = h, \quad \angle PDH = x, \quad \angle QDH = y, \quad k(O, R).$$

We have

$$x, y \in (-90^\circ, 90^\circ).$$

If $x = -y$, then $PQP'Q'$ is an isosceles trapezoid and the condition that the center of its circumcircle lies on k is equivalent to

$$PQP'Q' \text{ is a rectangle} \Leftrightarrow PQ = P'Q'.$$

Now let $x + y \neq 0^\circ$. Then if the center of the circumcircle of $PQP'Q'$ lies on k , we have that the perpendicular bisectors of PQ and $P'Q'$ intersect on k . Let that point be T . From the first bisector, T 's projection M onto l must be the midpoint of PQ , hence $HM = h(\tan x + \tan y)/2$ (the distance being signed and depending on the signs of x and y). From the second bisector, it follows that T is the midpoint of either the major or minor arc of $P'Q'$, so it forms with OD angles $x + y$ or $180^\circ + x + y$ around O . Therefore, its distance from OH is $\pm R \sin(x + y)$, depending on which arc midpoint T is. Therefore

$$\begin{aligned} \frac{1}{2}h(\tan x + \tan y) &= \pm R \sin(x + y) \Leftrightarrow \\ \frac{h}{\cos x \cos y} &= \pm 2R \Leftrightarrow \\ \frac{1}{2}h(\tan x - \tan y) &= \pm R \sin(x - y) \Leftrightarrow \\ P'Q' &= \pm PQ, \end{aligned}$$

where exactly one of the $+$ or $-$ must be true.

The converse direction does not need consideration of cases, since $\sin(x - y) \neq 0$ and therefore the argument flows backward.

4.3

The excircles constructed on the sides AB , BC , CA of $\triangle ABC$ touch the lines CA , CB , AB , AC , BC , BA respectively at the points C_1 , C_2 , A_1 , A_2 , B_1 , B_2 . Let

$$P_C = B_1B_2 \cap A_1A_2, \quad P_A = B_1B_2 \cap C_1C_2, \quad P_B = A_1A_2 \cap C_1C_2.$$

Prove that $P_C C$, $P_A A$, $P_B B$ pass through the center of the circumcircle of $\triangle P_A P_B P_C$.

Solution: Let the radius of the circumcircle of $\triangle ABC$ be $R = 1/2$, and the angles of the triangle be 2α , 2β , 2γ . Let A_1A_2 intersect the altitude through C at point P' , and let the foot of this altitude be C_H . We have


$$\begin{aligned} P'C_H &= C_H A_1 \cot \alpha = (AA_1 - AC_H) \cot \alpha \\ &= (2 \cos \alpha \cos \beta \cos \gamma - \sin 2\beta \cos 2\alpha) \cot \alpha \\ &= 2 \cos \beta (\cos \alpha \sin(\alpha + \beta) - \sin \beta \cos 2\alpha) \cot \alpha \\ &= 2 \cos \beta (\sin \alpha \cos \alpha \cos \beta + \sin \beta \sin^2 \alpha) \cot \alpha \\ &= 2 \cos \beta \sin \alpha \cos \alpha \cos(\alpha - \beta) \cot \alpha \\ &= 2 \cos \beta \cos \alpha \cos(\alpha - \beta) \end{aligned}$$

which is a symmetric expression in α and β . Then B_1B_2 intersects the altitude CC_H in the same point P' , i.e. $P \equiv P_C$. Therefore, $P_C C \perp AB$ and $H \in P_C C$. Analogously $H \in P_B B$ and $H \in P_A A$.

We shall show that H is the circumcenter of $\triangle P_A P_B P_C$. We have

$$\begin{aligned} P_C H &= 2 \cos \alpha \cos \beta \cos(\alpha - \beta) - \cos(2\alpha) \cos(2\beta) \\ &= (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \cos(\alpha - \beta) - \frac{\cos(2\alpha - 2\beta) + \cos(2\alpha + 2\beta)}{2} \\ &= \cos^2(\alpha - \beta) + \cos(\alpha - \beta) \cos(\alpha + \beta) - \frac{2 \cos^2(\alpha - \beta) - 1}{2} + \frac{\cos(2\gamma)}{2} \\ &= \frac{1 + \cos(2\alpha) + \cos(2\beta) + \cos(2\gamma)}{2}, \end{aligned}$$

which is a symmetric expression in α, β, γ , so we conclude that $P_C H = P_A H = P_B H$ and the statement is proved. \square



4.4

Let P be a point inside the acute triangle ABC . X is such a point that the quadrilateral $PBXC$ is inscribed in a circle and is with perpendicular diagonals. Similarly the points Y, Z are defined with respect to CA and AB . Prove that if $PB \perp XZ$ and $PA \perp YZ$, then P is the center of the circumcircle of $\triangle ABC$.

Solution: Let

$$M = PX \cap BC, \quad N = PY \cap AC, \quad L = PZ \cap AB.$$

Let the lengths of the sides of the triangle be $2a, 2b, 2c$, and let x, y, z be the directed distances (clockwise) from the midpoints on BC, CA, AB to M, N, L respectively.

From ABC acute and P interior it follows that

$$-1 < \frac{x}{a}, \frac{y}{b}, \frac{z}{c} < 1,$$

so

$$a \pm x > 0, b \pm y > 0, c \pm z > 0. \quad (a)$$

From $PB \perp XZ$ and Pythagorean theorem to $\triangle PBZ$ and $\triangle PBX$ it follows that

$$\begin{aligned} BZ^2 - BX^2 &= PZ^2 - PX^2 \\ \Rightarrow BL^2 + LZ^2 - BM^2 - MX^2 &= (PL + LZ)^2 - (PM + MX)^2 \\ &= PL^2 + LZ^2 + 2PL \cdot LZ \\ &\quad - PM^2 - MX^2 - 2PM \cdot MX \\ \Rightarrow BL^2 - BM^2 &= (PB^2 - BL^2) + 2PL \cdot LZ \\ &\quad - (PB^2 - BM^2) - 2PM \cdot MX \\ &= -BL^2 + 2BL \cdot LA + BM^2 - 2BM \cdot MC \end{aligned}$$

since $PL \cdot LZ = BL \cdot LA$ and $PM \cdot MX = BM \cdot MC$. From here:

$$\begin{aligned} BL^2 - BM^2 &= BL \cdot LA - BM \cdot MC \\ \Rightarrow BL(BL - AL) &= BM(BM - MC) \\ \Rightarrow (c - z)2z &= -(a + x)2x \\ \Rightarrow x(x + a) &= z(z - c). \quad (*) \end{aligned}$$

Analogously we obtain

$$z(c + z) = y(b - y). \quad (**)$$

From

$$\begin{aligned} 0 &= PA^2 - PB^2 + PB^2 - PC^2 + PC^2 - PA^2 \\ &= AL^2 - LB^2 + BM^2 - MC^2 + CN^2 - NA^2 \\ &\Rightarrow ax + by + cz = 0. \end{aligned}$$

Using that and adding the equalities (*) and (**), we obtain

$$\begin{aligned} ax + x^2 + z^2 + cz &= z^2 - cz + y^2 - by \\ \Rightarrow x^2 + cz &= y^2 = y^2 + ax + by + cz \\ \Rightarrow y(b + y) &= x(x - a), \quad (***) \end{aligned}$$

i.e. the third equality of the form (*) is true, which also proves that $PC \perp YX$. Now, multiply the three equalities to get:


$$\begin{aligned} xyz(a + x)(b + y)(c + z) &= xyz(x - a)(y - b)(z - c) \\ \Rightarrow xyz((a + x)(b + y)(c + z) + (a - x)(b - y)(c - z)) &= 0. \end{aligned}$$

Since the expressions in parentheses are sums of positive numbers due to (a), it follows that $xyz = 0$, so one of x, y, z is 0. Without loss of generality, let $x = 0$.

Then from (***) and (*) with $x = 0$ we get

$$y(b + y) = 0, \quad z(z - c) = 0,$$

which, owing to (a), implies $y = z = 0$. That means that P projects to the midpoints of the sides of the triangle, hence it is the center of its circumcircle.



4.5

Let C_1, A_1, B_1 be the touchpoints on the incircle $\Gamma(I, r)$ with the sides AB, BC, CA of $\triangle ABC$ respectively. Also let $\angle A = 60^\circ$ and $A_2 = \Gamma \cap AA_1$, $A_2 \neq A_1$. If $I \in BC$ and A, A_2, I, T lie on a circle, find $\angle TA_1C_1$.

Solution: If there were two distinct points T_1, T_2 on B_1C_1 such that A, A_2, T_1, T_2, I lie on a circle, then we would have

$$\angle AT_1I + \angle AT_2I = 180^\circ,$$

which would be impossible since both $\angle AT_1I$ and $\angle AT_2I$ would need to be obtuse (being larger than the rectangles $\angle AB_1I$ and $\angle AC_1I$). Thus the point in question is exactly one, and we will construct it.

Let M be the midpoint of the arc $B_1A_1C_1$ in Γ , and $T' = MA_2 \cap BC$, then A_2T' is the angular bisector of $\angle B_1A_2C_1$ and hence

$$\frac{B_1T'}{T'C_1} = \frac{B_1A_2}{A_2C_1} = \frac{\frac{B_1A_1}{AA_1} AB_1}{\frac{A_1C_1}{AA_1} AC_1} = \frac{B_1A_1}{A_1C_1}$$

Therefore A_1T' is the angular bisector of $\angle B_1A_1C_1$ and $\angle T'A_1C_1 = 30^\circ$.


We shall show that $T' \equiv T$ and the problem is solved. We have

$$\begin{aligned} \angle B_1A_2M &= \angle MA_2C_1 = \angle T'B_1M \Rightarrow \\ \triangle MB_1A_2 &\sim \triangle MTB_1 \Rightarrow \\ \frac{MB_1}{MT'} &= \frac{MA_2}{MB_1} \Rightarrow MB_1^2 = MT' \cdot MA_2. \end{aligned}$$

Since $I \in AM$, then from $\angle IB_1M = \angle IMB_1 = 30^\circ = \angle B_1AM$ it follows

$$\triangle MB_1I \sim \triangle MAB_1 \Rightarrow MI \cdot MA = MB_1^2 = MT' \cdot MA_2$$

and therefore A, A_2, I, T' lie on a circle, from where $T' \equiv T$. Final answer: 30° □




4.6

Given is $\triangle ABC$ with $\angle B = 60^\circ$. Let T be the point of tangency of the incircle of $\triangle ABC$ with AB , and C' be such that T is the midpoint of CC' . If the perpendicular bisector of BC' intersects the angle bisector of $\angle A$ at A' , prove that $\triangle A'BC'$ is equilateral.

Solution: Let I be the center of the inscribed circle in $\triangle ABC$, and let it touch BC at T' . Let $L = AI \cap BC$ and $M = CI \cap AB$. Then we shall show that $BL + BM = 2BT$. Indeed, $\angle IMB + \angle ILB = \angle A + \frac{1}{2}\angle C + \angle C + \frac{1}{2}\angle A = \frac{3}{2}(\angle A + \angle C) = 180^\circ$ since $\angle A + \angle C = 180^\circ - \angle B = 120^\circ$. Furthermore, $BM > BT \iff BL < BT' = BT$. Then the right triangles $\triangle IMT$ and $\triangle ILT'$ are identical, since they have the same angles and since $IT = IT'$ (being the radius of the incircle). Now from $MT = LT'$ it follows that $BL + BM = 2BT$.

Let N be obtained from L by rotation by 60° about B , i.e. $N \in AB$ and $BN = BL$. Then $MT = TN$ and $CMCN$ is a parallelogram. So $\angle BNC = 180^\circ - \angle BMC = \angle BLA$. Then under rotation by -60° about B the ray NC' goes into LA , hence C' goes into a point B' on LA . It is easy to see that $\triangle BC'B'$ is equilateral, hence B' lies on the perpendicular bisector of BC' . That means $B' = A'$ and so $\triangle A'BC'$ is equilateral.



4.7

Let Γ be a set of points A_1, A_2, \dots, A_n , $n > 2$, which is symmetric with respect to point O . If no three points of Γ are collinear and $OA_1 + OA_2 + \dots + OA_n > 7$, prove that we can choose two points P and Q from Γ , such that the circumradius of $\triangle OPQ$ be larger than 1.

Solution: From the condition for noncollinearity $\Rightarrow O \notin \Gamma$ and so $n = 2m$, $m > 1$. Without loss of generality, we may consider that $\arg(a_i) < \arg(a_j)$ for $i < j$, where a_i is the affix of A_i , and 0 is the affix of O . We are given that $a_i + a_{i+m} = 0$, $i = 1, 2, \dots, m$.

Let us consider the convex polygon $b = B_1B_2 \dots B_{2m}$ such that $b_i = a_1 + a_2 + \dots + a_i$. That it is convex follows from the fact that its interior angles are $< 180^\circ$ (between a_i and a_{i+m} there is a point a_{i+1} , since $m > 1$). Its sides have lengths OA_1, OA_2, \dots, OA_n and therefore its perimeter p is larger than 7. We shall prove the following two lemmas:

Lemma 1: Every circle containing b has perimeter $> p$.

Lemma 2: $\exists i \in \mathbb{N}$: the circumcircle of $\triangle B_iB_{i+1}B_{i+2}$ contains the whole polygon b ($B_{2m+1} = B_1$, $B_{2m+2} = B_2$).

From the two lemmas it follows that there exists a circumcircle of $\triangle B_iB_{i+1}B_{i+2}$ with radius r :

$$2\pi r > p > 7 \Rightarrow r > 1.$$


It remains to note that the radius of the circumcircle of $\triangle OA_iA_{i+1}$ is the same r and the problem is solved. Here are the proofs of the lemmas:

Lemma 1: Let us construct for every i from B_i the rays B_iC_i and B_iD_i outside b :

$$B_iC_i \perp B_iB_{i-1}, \quad B_iD_i \perp B_iB_{i+1}.$$

In the region $D_iB_iB_{i+1}C_{i+1}$ are contained arcs from the circle about b and each of them has length $> B_iB_{i+1}$ which proves the lemma.

Lemma 2: Let us take the side B_1B_2 and find the largest angle $\angle B_iB_{i+1}B_{i+2}$ ($n \geq i \geq 3$) among $\angle B_1B_2B_3$. Clearly the circle circumscribed about $\triangle B_iB_{i+1}B_{i+2}$ covers all points of b . Let us consider all triples $\{i, p, q\}$ ($0 < p < q < n$, $1 \leq i \leq n$) and take that one of them for which q is minimal and the circle circumscribed about $\triangle B_iB_{i+p}B_{i+q}$ contains b . Thus q is minimal and all triples are of the type $\{i, 1, i-1\}$ are one of them. Then if $q > 2 \Rightarrow p > 1$ or $q > p+1$. The two cases are analogous \Rightarrow if $p > 1$ we take the largest angle between $\angle B_iB_{i+p}B_{i+p+1}$ and the circle about $\triangle B_iB_{i+p}B_{i+p+1}$ contains b . But then $p < q \Rightarrow$ the triple $\{i, i-1, p\}$ is more optimal, which is impossible. Finally $q = 2$, $p = 1$ and the lemma is proved.



4.8

Let P and Q be points inside an acute triangle $\triangle ABC$ such that

$$\angle PCA = \angle PAB = \angle PBC, \quad \angle QBA = \angle QCB = \angle QAC.$$

- (a) Prove that the feet of the perpendiculars from P and Q to the sides of $\triangle ABC$ lie on a circle whose center is the midpoint of PQ .
- (b) If O is the circumcenter of $\triangle ABC$, prove that $OP = OQ$.

Solution (idea of G. Panova): Let $\triangle A_1B_1C_1$ denote the pedal triangle of P with respect to $\triangle ABC$, where $A_1 \in BC$, $B_1 \in CA$, $C_1 \in AB$. By the sine law, we have

$$\frac{\sin \angle PAC}{\sin \angle PCA} \cdot \frac{\sin \angle PCB}{\sin \angle PBC} \cdot \frac{\sin \angle PBA}{\sin \angle PAB} = 1 = \frac{\sin \angle QCA}{\sin \angle QAC} \cdot \frac{\sin \angle QBC}{\sin \angle QCB} \cdot \frac{\sin \angle QAB}{\sin \angle QBA}.$$

Let $x = \angle PCA = \angle PAB = \angle PBC$ and $y = \angle QBA = \angle QCB = \angle QAC$. Also, let $\alpha = \angle BAC$, $\beta = \angle CBA$, $\gamma = \angle ACB < 90^\circ$ be the angles of $\triangle ABC$. Then x and y are two solutions to the equation $f(x) = 0$, where

$$f(x) = \sin^3 x - \sin(\alpha - x) \sin(\beta - x) \sin(\gamma - x). \quad (*)$$

If $\mu = \min(\alpha, \beta, \gamma)$, then $f(x)$ is a continuous, strictly increasing function on $(0, \mu) \subset (0, 90^\circ)$ and its values range from $-\sin \alpha \sin \beta \sin \gamma$ to $\sin^3 \mu$. Therefore f has a unique root, so $x = y$.

Observe that the quadrilaterals PB_1AC_1 and PB_1CA_1 are cyclic, since each quadrilateral has two opposite right angles. Then

$$\angle PB_1C_1 = \angle PAC_1 = x, \quad \angle PB_1A_1 = \angle PCB = \gamma - x,$$

hence $\angle C_1B_1A_1 = \gamma = \angle ACB$. Similarly, $\angle A_1C_1B_1 = \angle BAC = \alpha$, so the triangles $\triangle ABC$ and $\triangle C_1A_1B_1$ are similar and similarly oriented. Then there exists a spiral similarity ρ_1 centered at P mapping $A_1 \rightarrow B$, $B_1 \rightarrow C$, $C_1 \rightarrow A$ respectively. Since $\rho_1(B_1) = C$, we see that the angle of ρ_1 is $\angle B_1PC = 90^\circ - x$ and its coefficient is $CP/B_1P = 1/\sin x$.

Analogously, let $\triangle A_2B_2C_2$ be the pedal triangle of Q with respect to $\triangle ABC$, where $A_2 \in BC$, $B_2 \in CA$, $C_2 \in AB$. By the earlier argument, there exists a spiral similarity ρ_2 with center Q , angle $90^\circ - x$ and coefficient $\sin(x)$, mapping $\triangle ABC$ to $\triangle B_2C_2A_2$.

Since the coefficient of $\rho_2\rho_1$ equals the product of the coefficients of ρ_1 and ρ_2 , which is 1, it follows that $\rho = \rho_2\rho_1$ is a rotation that maps $\triangle C_1A_1B_1$ to $\triangle B_2C_2A_2$.

Let O' be the midpoint of PQ and $S = \rho_1(O')$. Since ρ_1 has center P , coefficient $1/\sin(x)$, and angle $90^\circ - x$, we have $PS = PO'/\sin(x)$ and $\angle O'PS = 90^\circ - x$. If K is the projection of S onto PQ , we have $\angle O'PS = 90^\circ - x$, so

$$PK = PS \cos(90^\circ - x) = PS \sin x = PO'.$$

Hence $K = O'$, so $SO' \perp PQ$. Therefore S lies on the perpendicular bisector of PQ , and thus $PS = QS$. Therefore $\rho_2(S) = O'$, consequently $\rho(O') = O'$ and O' is the center of ρ . Since

$$\rho(C_1) = B_2, \quad \rho(A_1) = C_2, \quad \rho(B_1) = A_2,$$

then

$$O'C_1 = O'B_2, \quad O'A_1 = O'C_2, \quad O'B_1 = O'A_2. \quad (**)$$

At the same time, since $PC_1 \perp AB$ and $QC_2 \perp AB$, both P and Q project onto AB at C_1 and C_2 respectively. Hence the midpoint O' of PQ lies on the perpendicular bisector of C_1C_2 and so $O'C_1 = O'C_2$. Analogously, $O'A_1 = O'A_2$. Combined with (**), that yields

$$O'B_2 = O'C_1 = O'C_2 = O'A_1 = O'A_2 = O'B_1,$$

hence O' is the circumcenter of the hexagon $C_1A_1B_2C_2B_1A_2$ and (a) is proved.

To prove (b), let us note that since O' is the circumcenter of $\triangle C_1A_1B_1$, its image under ρ_1 is the circumcenter of $\triangle ABC$, hence $S = \rho_1(O') = O$. From $PS = QS$ we obtain $PO = QO$ and the problem is solved. \square

